arXiv:0806.3112v2 [math.QA] 24 Oct 2008

Mirković-Vilonen polytopes lying in a Demazure crystal and an opposite Demazure crystal

Satoshi Naito
Institute of Mathematics, University of Tsukuba,
Tsukuba, Ibaraki 305-8571, Japan (e-mail: `naito@math.tsukuba.ac.jp`)

and

Daisuke Sagaki
Institute of Mathematics, University of Tsukuba,
Tsukuba, Ibaraki 305-8571, Japan (e-mail: `sagaki@math.tsukuba.ac.jp`)

Abstract

We give a necessary and sufficient condition for an MV polytope P in a highest weight crystal to lie in an arbitrary fixed Demazure crystal (resp., opposite Demazure crystal), in terms of the lengths of edges along a path through the 1-skeleton of P corresponding to a reduced word for the longest element of the Weyl group W. Also, we give an explicit description as a pseudo-Weyl polytope for extremal MV polytopes in a highest weight crystal. Finally, by combining the results above, we obtain a polytopal condition for an MV polytope P to lie in an arbitrary fixed opposite Demazure crystal.

1 Introduction.

Let G be a complex, connected, semisimple algebraic group with Lie algebra $\mathfrak{g}$, T a maximal torus with Lie algebra $\mathfrak{h}$, B a Borel subgroup containing T, N the unipotent radical of B, and $G^\vee$ the Langlands dual group of G with Lie algebra $\mathfrak{g}^\vee$; we choose the convention that the roots in B are the negative roots. Throughout this paper, we assume that G is simply-laced; all highest weight crystals and Demazure crystals — the ones of our interest — can be obtained from those in the simply-laced cases by a standard technique of "folding" by diagram automorphisms (see [NS1] – [NS3], and [Ho]). The affine Grassmannian $\mathcal{G}r$ for G is defined to be the quotient set $G(\mathcal{K})/G(\mathcal{O})$, equipped with the structure of a complex algebraic ind-variety, where $G(\mathcal{K})$ denotes the group of $\mathcal{K} := \mathbb{C}(\!(t)\!)$-valued points of G, and $G(\mathcal{O}) \subset G(\mathcal{K})$ denotes the subgroup of $\mathcal{O} := \mathbb{C}[[t]]$-valued points of G. For each dominant coweight $\lambda \in X_*(T) := \mathrm{Hom}(\mathbb{C}^*, T)$, Mirković and Vilonen [MiV1], [MiV2] discovered a family $\mathcal{Z}(\lambda)$ of closed, irreducible, algebraic subvarieties, called Mirković-Vilonen (MV for short) cycles, in the closure $\overline{\mathcal{G}r^\lambda} \subset \mathcal{G}r$ of the $G(\mathcal{O})$-orbit $\mathcal{G}r^\lambda := G(\mathcal{O})[\lambda] \subset \mathcal{G}r$ through the image $[\lambda] \in \mathcal{G}r$ of $\lambda \in X_*(T) \subset G(\mathcal{K})$, and proved that these algebraic subvarieties provide a basis for the irreducible highest weight module $L(\lambda)$ of highest weight λ over the Langlands dual group $G^\vee$; more precisely, the family $\mathcal{Z}(\lambda)$ consists of the irreducible components of the closure of the intersection $\mathcal{G}r^\lambda \cap N(\mathcal{K})[\nu] \subset \mathcal{G}r$ for weights $\nu \in X_*(T)$ of $L(\lambda)$, where $N(\mathcal{K}) \subset G(\mathcal{K})$ denotes the subgroup of $\mathcal{K}$-valued points of G. Furthermore, Braverman and Gaitsgory [BrG] (see also [BFG]) endowed the set $\mathcal{Z}(\lambda)$ of MV cycles with a crystal structure, and gave an isomorphism of crystals between the resulting crystal and the crystal basis $\mathcal{B}(\lambda)$ of the irreducible highest weight module $V(\lambda)$ of highest weight $\lambda \in X_*(T) \subset \mathfrak{h}$ for the quantized universal enveloping algebra $U_q(\mathfrak{g}^\vee)$ over $\mathbb{C}(q)$ of the (Langlands) dual Lie algebra $\mathfrak{g}^\vee$ of $\mathfrak{g}$ with Cartan subalgebra $\mathfrak{h}^*$.

In order to obtain an explicit combinatorial description of the MV cycles in $\mathcal{Z}(\lambda)$ for a dominant coweight $\lambda \in X_*(T) \subset \mathfrak{h}$, Anderson [An] defined a family $\mathcal{MV}(\lambda)$ of convex polytopes in the real form $\mathfrak{h}_\mathbb{R}$ of $\mathfrak{h}$, which he called Mirković-Vilonen (MV for short) polytopes, to be the moment map images of the MV cycles in $\mathcal{Z}(\lambda)$. Furthermore, Kamnitzer [Kam1], [Kam2] characterized the MV polytopes in $\mathcal{MV}(\lambda)$ as certain pseudo-Weyl polytopes with highest vertex λ satisfying the "tropical Plücker relations", and then showed the existence of a bijection (in fact, an isomorphism of crystals) between the crystal basis $\mathcal{B}(\lambda)$ and the set $\mathcal{MV}(\lambda)$ of MV polytopes, which is endowed with a crystal structure coming from the one (due to Lusztig [Lu1] and Berenstein-Zelevinsky [BeZ]) on the canonical basis for the negative part $U_q^-(\mathfrak{g}^\vee)$ of $U_q(\mathfrak{g}^\vee)$; in [Kam2], Kamnitzer also proved that this crystal structure agrees with the crystal structure coming from the one on the set $\mathcal{Z}(\lambda)$ of MV cycles.

Let W denote the Weyl group of $\mathfrak{g}$, with e the unit element and w_0 the longest element. We fix a dominant coweight $\lambda \in X_*(T) \subset \mathfrak{h}_\mathbb{R}$. Now, to understand better the MV polytopes in $\mathcal{MV}(\lambda)$, we consider a filtration (compatible with the Bruhat ordering $\leq$ on W) of the crystal basis $\mathcal{B}(\lambda)$ by Demazure crystals $\mathcal{B}_x(\lambda) \subset \mathcal{B}(\lambda)$ (resp., opposite Demazure crystals $\mathcal{B}^x(\lambda) \subset \mathcal{B}(\lambda)$), $x \in W$, and then their images $\mathcal{MV}_x(\lambda) \subset \mathcal{MV}(\lambda)$ (resp., $\mathcal{MV}^x(\lambda) \subset \mathcal{MV}(\lambda)$), $x \in W$, under the isomorphism of crystals $\mathcal{B}(\lambda) \xrightarrow{\sim} \mathcal{MV}(\lambda)$ above. Here, for each $x \in W$, the Demazure module $V_x(\lambda)$ (resp., opposite Demazure module $V^x(\lambda)$) is the $U_q^+(\mathfrak{g}^\vee)$-submodule (resp., $U_q^-(\mathfrak{g}^\vee)$-submodule) of $V(\lambda)$ generated by the one-dimensional weight space $V(\lambda)_{x\cdot\lambda} \subset V(\lambda)$ of weight $x \cdot \lambda \in X_*(T) \subset \mathfrak{h}_\mathbb{R}$, where $U_q^+(\mathfrak{g}^\vee)$ (resp., $U_q^-(\mathfrak{g}^\vee)$) is the positive part (resp., negative part) of $U_q(\mathfrak{g}^\vee)$; recall from [Kas2] that the Demazure crystal $\mathcal{B}_x(\lambda)$ (resp., opposite Demazure crystal $\mathcal{B}^x(\lambda)$) is a subset of $\mathcal{B}(\lambda)$ such that

$$V(\lambda) \supset V_x(\lambda) = \bigoplus_{b \in \mathcal{B}_x(\lambda)} \mathbb{C}(q) G_\lambda(b) \qquad \left(\text{resp., } V(\lambda) \supset V^x(\lambda) = \bigoplus_{b \in \mathcal{B}^x(\lambda)} \mathbb{C}(q) G_\lambda(b) \right),$$

where $G_\lambda(b)$, $b \in \mathcal{B}(\lambda)$, form the lower global basis of $V(\lambda)$. In this paper, for a dominant coweight $\lambda \in X_*(T) \subset \mathfrak{h}_\mathbb{R}$ and $x \in W$, we give a necessary and sufficient condition for an MV polytope in $\mathcal{MV}(\lambda)$ to lie in the image $\mathcal{MV}_x(\lambda) \subset \mathcal{MV}(\lambda)$ (resp., $\mathcal{MV}^x(\lambda) \subset \mathcal{MV}(\lambda)$)

of $\mathcal{B}_x(\lambda) \subset \mathcal{B}(\lambda)$ (resp., $\mathcal{B}^x(\lambda) \subset \mathcal{B}(\lambda)$) under the isomorphism $\mathcal{B}(\lambda) \xrightarrow{\sim} \mathcal{MV}(\lambda)$, in terms of the lengths of edges along a path through the 1-skeleton of an MV polytope corresponding to a reduced word of $w_0 \in W$; see Theorem 3.2.1 (1) (resp., Theorem 3.5.1) for details. Also, we give a similar result for the crystal basis $\mathcal{B}(\infty)$ of the negative part $U_q^-(\mathfrak{g}^\vee)$ of $U_q(\mathfrak{g}^\vee)$; see Theorem 3.2.1 (2) for details.

Furthermore, for each $x \in W$, we give an explicit description as a pseudo-Weyl polytope for the image (called an extremal MV polytope) $P_{x\cdot\lambda} \in \mathcal{MV}(\lambda)$ of the extremal element $u_{x\cdot\lambda} \in \mathcal{B}(\lambda)$ of weight $x\cdot\lambda$ under the isomorphism of crystals $\mathcal{B}(\lambda) \xrightarrow{\sim} \mathcal{MV}(\lambda)$; see Theorem 4.1.5 (1) for details. As a corollary, we obtain a purely combinatorial proof of the fact that the extremal MV polytope $P_{x\cdot\lambda} \in \mathcal{MV}(\lambda)$ of weight $x\cdot\lambda \in X_*(T) \subset \mathfrak{h}_\mathbb{R}$ is identical to the convex hull $\mathrm{Conv}(W_{\le x}\cdot\lambda)$ in $\mathfrak{h}_\mathbb{R}$ of the subset $W_{\le x}\cdot\lambda := \bigl\{z\cdot\lambda \mid z \in W \text{ with } z \le x\bigr\}$ of the W-orbit $W\cdot\lambda$ through λ; see Theorem 4.1.5 (2) for details. This result is a generalization of the fact (see [An] and [Kam1]) that the MV polytope $P_{e\cdot\lambda} \in \mathcal{MV}(\lambda)$ of highest weight λ consists only of the single element $\lambda \in X_*(T) \subset \mathfrak{h}$, which is the moment map image of the point $[\lambda] \in \mathcal{G}r$, and the MV polytope $P_{w_0\cdot\lambda} \in \mathcal{MV}(\lambda)$ of lowest weight $w_0\cdot\lambda$ is identical to the convex hull $\mathrm{Conv}(W\cdot\lambda)$ in $\mathfrak{h}_\mathbb{R}$ of the W-orbit $W\cdot\lambda$, which is the moment map image of the finite-dimensional projective algebraic subvariety $\overline{\mathcal{G}r^\lambda}$ of $\mathcal{G}r$.

By combining our result for opposite Demazure crystals and the explicit description above of extremal MV polytopes, we can prove that for a dominant coweight $\lambda \in X_*(T) \subset \mathfrak{h}_\mathbb{R}$ and $x \in W$, an MV polytope P in $\mathcal{MV}(\lambda)$ lies in the opposite Demazure crystal $\mathcal{MV}^x(\lambda)$ if and only if P contains (as a set) the extremal MV polytope $P_{x\cdot\lambda} = \mathrm{Conv}(W_{\le x}\cdot\lambda)$; see Theorem 4.5.1.

Finally, for a dominant coweight $\lambda \in X_*(T) \subset \mathfrak{h}_\mathbb{R}$, we refine the filtration of the crystal $\mathcal{MV}(\lambda)$ consisting of Demazure crystals $\mathcal{MV}_x(\lambda)$ (resp., opposite Demazure crystals $\mathcal{MV}^x(\lambda)$), $x \in W$, to obtain the following decomposition into a disjoint union:

$$\mathcal{MV}(\lambda) = \bigsqcup_{z \in W^\lambda_{\min}} \overline{\mathcal{MV}}_z(\lambda) \qquad \left(\text{resp., } \mathcal{MV}(\lambda) = \bigsqcup_{z \in W^\lambda_{\max}} \overline{\mathcal{MV}}^z(\lambda)\right),$$

where $W^\lambda_{\min}$ (resp., $W^\lambda_{\max}$) is the set of minimal (resp., maximal) coset representatives for the quotient set W/W_λ, with W_λ ($\subset W$) the stabilizer of λ, with respect to the Bruhat ordering $\le$ on W, and where, for each $x \in W^\lambda_{\min}$ (resp., $x \in W^\lambda_{\max}$), we set

$$\overline{\mathcal{MV}}_x(\lambda) := \mathcal{MV}_x(\lambda) \setminus \left(\bigcup_{z \in W^\lambda_{\min},\ z < x} \mathcal{MV}_z(\lambda) \right)$$

$$\left(\text{resp., } \quad \overline{\mathcal{MV}}^x(\lambda) := \mathcal{MV}^x(\lambda) \setminus \left(\bigcup_{z \in W^\lambda_{\max},\ z > x} \mathcal{MV}^z(\lambda) \right) \right).$$

Then, for each $x \in W$, we give a necessary and sufficient condition for an MV polytope $P \in \mathcal{MV}(\lambda)$ to lie in $\overline{\mathcal{MV}}_x(\lambda)$ (resp., $\overline{\mathcal{MV}}^x(\lambda)$); see Proposition 5.1.4 (resp., Proposition 5.2.5) for details. This result tells us the smallest, with respect to the inclusion relation, Demazure crystal (resp., opposite Demazure crystal) in which $P \in \mathcal{MV}(\lambda)$ lies.

Here, we should mention some closely related results: in [Sc, §11.2], Schwer showed that the dimension of each weight space of a Demazure module is given by the number of top-dimensional irreducible components of the intersection of an $N(\mathcal{K})$-orbit and an orbit under the standard Iwahori subgroup of $G(\mathcal{K})$ in the affine Grassmannian $\mathcal{G}r$ (note that the closures of these irreducible components are not MV cycles) ; in [I, §4], Ion showed a very similar result by using certain intersections in the affine (full) flag manifold; in [Sa], Savage gave a geometric realization of Demazure crystals for simply-laced Kac-Moody algebras by using certain subvarieties (called Demazure quiver varieties) of Nakajima's quiver varieties; in [N], Nakashima gave a condition for an element of a highest weight crystal to lie in a Demazure crystal, and also gave an explicit description of extremal elements for symmetrizable Kac-Moody algebras, both of which are in terms of polyhedral realizations; among other realizations of Demazure crystals, we name [Li1], [Li3], and [LP].

This paper is organized as follows. In §§2.1 and 2.2, following Kamnitzer [Kam1], [Kam2], we review some of the basic facts on MV polytopes, and then give a description of the crystal structure on them. In §§3.1 and 3.2, after recalling the notion of Demazure crystals, we state a necessary and sufficient condition for an MV polytope to lie in a Demazure crystal in terms of the lengths of its edges; the proof is given in §3.3. In §§3.4 and 3.5, a similar result for an opposite Demazure crystal is obtained. In §4.1, we provide an explicit description of extremal MV polytopes; §§4.2–4.4 are devoted to its proof. In §4.5, we give a polytopal condition for an MV polytope to lie in an opposite Demazure crystal, and in §4.6, we pose a question concerning a necessary (but, not sufficient) polytopal condition for an MV polytope to lie in a Demazure crystal. Finally, in §5, we give a necessary and sufficient condition for an MV polytope to lie in the $\overline{\mathcal{MV}}_x(\lambda)$ (resp., $\overline{\mathcal{MV}}^x(\lambda)$) above for $x \in W$.

2 Mirković-Vilonen polytopes.

2.1 Mirković-Vilonen polytopes. Here, following [Kam1], we briefly review some of the basic facts on Gelfand-Goresky-MacPherson-Serganova (GGMS for short) data and Mirković-Vilonen (MV for short) polytopes. Let G be a complex, connected, semisimple algebraic group with Lie algebra $\mathfrak{g}$, T a maximal torus with Lie algebra $\mathfrak{h}$, B a Borel subgroup containing T, and N the unipotent radical of B; we choose the convention that the roots in B are the negative roots. Throughout this paper, we assume that G (and hence $\mathfrak{g}$) is simply-laced, for reasons explained in the Introduction. Let $\bigl(A = (a_{ij})_{i,j \in I}, \Pi := \{\alpha_j\}_{j \in I}, \Pi^\vee := \{h_j\}_{j \in I}, \mathfrak{h}^*, \mathfrak{h}\bigr)$ be the root datum of $\mathfrak{g}$, where $A = (a_{ij})_{i,j \in I}$ is the Cartan matrix, $\mathfrak{h}$ is the Cartan subalgebra, $\Pi := \{\alpha_j\}_{j \in I} \subset \mathfrak{h}^* := \mathrm{Hom}_{\mathbb{C}}(\mathfrak{h}, \mathbb{C})$ is the set of simple roots, and $\Pi^\vee := \{h_j\}_{j \in I} \subset \mathfrak{h}$ is the set of simple coroots; note that $\langle \alpha_j, h_i \rangle = a_{ij}$ for $i, j \in I$, where $\langle \cdot, \cdot \rangle$ denotes the canonical pairing between $\mathfrak{h}^*$ and $\mathfrak{h}$. Let $W := \langle s_j \mid j \in I \rangle$ be the Weyl group of $\mathfrak{g}$, where s_j, $j \in I$, are the simple reflections, with length function $\ell : W \to \mathbb{Z}_{\geq 0}$, and let $e, w_0 \in W$ be the unit element and the longest element of W, respectively. Let $R(w_0)$ denote the set of all reduced words for w_0, that is, all sequences $(i_1, i_2, \ldots, i_m)$ of elements of I such that $s_{i_1} s_{i_2} \cdots s_{i_m} = w_0$, where m is the length $\ell(w_0)$ of the longest element w_0. Also, we denote by $\leq$ the (strong) Bruhat ordering on W.

We denote by $G^\vee$ the Langlands dual group of G, and by $\mathfrak{g}^\vee$ its Lie algebra. Thus, $\mathfrak{g}^\vee$ is the complex finite-dimensional semisimple Lie algebra associated to the root datum $\bigl({}^t A = (a_{ji})_{i,j \in I}, \Pi^\vee := \{h_j\}_{j \in I}, \Pi := \{\alpha_j\}_{j \in I}, \mathfrak{h}, \mathfrak{h}^*\bigr)$; note that the Cartan subalgebra of $\mathfrak{g}^\vee$ is $\mathfrak{h}^*$, not $\mathfrak{h}$. Let $U_q(\mathfrak{g}^\vee)$ denote the quantized universal enveloping algebra over the field $\mathbb{C}(q)$ of rational functions in q associated to the dual Lie algebra $\mathfrak{g}^\vee$, with positive part $U_q^+(\mathfrak{g}^\vee)$ and negative part $U_q^-(\mathfrak{g}^\vee)$. We denote by $\mathcal{B}(\infty)$ the crystal basis of $U_q^-(\mathfrak{g}^\vee)$. Also, for a dominant coweight $\lambda \in X_*(T) := \mathrm{Hom}(\mathbb{C}^*, T) \subset \mathfrak{h}_\mathbb{R} := \sum_{j \in I} \mathbb{R} h_j \subset \mathfrak{h}$, denote by $V(\lambda)$ the integrable highest weight $U_q(\mathfrak{g}^\vee)$-module of highest weight λ, and by $\mathcal{B}(\lambda)$ the crystal basis of $V(\lambda)$.

Let $\mu_\bullet = (\mu_w)_{w \in W}$ be a collection of elements of $\mathfrak{h}_\mathbb{R} = \sum_{j \in I} \mathbb{R} h_j$. We call $\mu_\bullet$ a Gelfand-Goresky-MacPherson-Serganova (GGMS) datum if it satisfies the condition that $w^{-1} \cdot (\mu_{w'} - \mu_w) \in \sum_{j \in I} \mathbb{Z}_{\geq 0} h_j$ for all $w, w' \in W$. It follows by induction on W with respect to the (weak) Bruhat ordering that $\mu_\bullet = (\mu_w)_{w \in W}$ is a GGMS datum if and only if

$$\mu_{ws_i} - \mu_w \in \mathbb{Z}_{\geq 0}\,(w \cdot h_i) \quad \text{for every } w \in W \text{ and } i \in I. \tag{2.1.1}$$

Following [Kam1] and [Kam2], to each GGMS datum $\mu_\bullet = (\mu_w)_{w \in W}$, we associate a convex polytope $P(\mu_\bullet) \subset \mathfrak{h}_\mathbb{R}$ by:

$$P(\mu_\bullet) = \bigl\{ h \in \mathfrak{h}_\mathbb{R} \mid w^{-1} \cdot (h - \mu_w) \in \textstyle\sum_{j \in I} \mathbb{R}_{\geq 0} h_j \text{ for all } w \in W \bigr\}; \tag{2.1.2}$$

the polytope $P(\mu_\bullet)$ is called a pseudo-Weyl polytope with GGMS datum $\mu_\bullet$. Note that the GGMS datum $\mu_\bullet = (\mu_w)_{w \in W}$ is determined uniquely by the convex polytope $P(\mu_\bullet)$. Also, we know from [Kam1, Proposition 2.2] that the set of vertices of the polytope $P(\mu_\bullet)$ is given by the collection $\mu_\bullet = (\mu_w)_{w \in W}$ (possibly, with repetitions). In particular, we have

$$P(\mu_\bullet) = \mathrm{Conv}\,\bigl\{ \mu_w \mid w \in W \bigr\},$$

where for a subset X of $\mathfrak{h}_\mathbb{R}$, $\mathrm{Conv}\, X$ denotes the convex hull in $\mathfrak{h}_\mathbb{R}$ of X.

Definition 2.1.1. Let $\mathbf{i} = (i_1, i_2, \ldots, i_m) \in R(w_0)$ and $\mathbf{j} = (j_1, j_2, \ldots, j_m) \in R(w_0)$ be reduced words for the longest element $w_0 \in W$.

(1) We say that $\mathbf{i}$ and $\mathbf{j}$ are related by a 2-move if there exist indices $i, j \in I$ with $a_{ij} = a_{ji} = 0$ and an integer $0 \leq k \leq m-2$ such that $i_l = j_l$ for all $1 \leq l \leq m$ with

$l \neq k+1,\ k+2$, and such that $i_{k+1} = j_{k+2} = i$, $i_{k+2} = j_{k+1} = j$, i.e.,

$$\begin{aligned} \mathbf{i} &= (i_1,\ \ldots,\ i_k,\ i,\ j,\ i_{k+3},\ \ldots,\ i_m), \\ \mathbf{j} &= (i_1,\ \ldots,\ i_k,\ j,\ i,\ i_{k+3},\ \ldots,\ i_m). \end{aligned} \tag{2.1.3}$$

(2) We say that $\mathbf{i}$ and $\mathbf{j}$ are related by a 3-move if there exist indices $i, j \in I$ with $a_{ij} = a_{ji} = -1$ and an integer $0 \le k \le m-3$ such that $i_l = j_l$ for all $1 \le l \le m$ with $l \neq k+1,\ k+2,\ k+3$, and such that $i_{k+1} = i_{k+3} = j_{k+2} = i$, $i_{k+2} = j_{k+1} = j_{k+3} = j$, i.e.,

$$\begin{aligned} \mathbf{i} &= (i_1,\ \ldots,\ i_k,\ i,\ j,\ i,\ i_{k+4},\ \ldots,\ i_m), \\ \mathbf{j} &= (i_1,\ \ldots,\ i_k,\ j,\ i,\ j,\ i_{k+4},\ \ldots,\ i_m). \end{aligned} \tag{2.1.4}$$

Remark 2.1.2. Let $\mathbf{i}, \mathbf{j} \in R(w_0)$. It is well-known (see, for example, [BjB, Theorem 3.3.1 (ii)]) that there exists a sequence $\mathbf{i} = \mathbf{i}_0,\ \mathbf{i}_1,\ \ldots,\ \mathbf{i}_t = \mathbf{j}$ of elements of $R(w_0)$ such that $\mathbf{i}_u$ and $\mathbf{i}_{u+1}$ are related by a 2-move or a 3-move for each $0 \le u \le t-1$.

Let $\mathbf{i} = (i_1,\ i_2,\ \ldots,\ i_m) \in R(w_0)$ be a reduced word for w_0. We set $w_l^{\mathbf{i}} := s_{i_1} s_{i_2} \cdots s_{i_l} \in W$ for $0 \le l \le m$. For a GGMS datum $\mu_\bullet = (\mu_w)_{w \in W}$, define integers $n_l^{\mathbf{i}} = n_l^{\mathbf{i}}(\mu_\bullet) \in \mathbb{Z}_{\ge 0}$, $1 \le l \le m$, via the following "length formula" (see [Kam1, Eq.(8)]):

$$\mu_{w_l^{\mathbf{i}}} - \mu_{w_{l-1}^{\mathbf{i}}} = n_l^{\mathbf{i}} w_{l-1}^{\mathbf{i}} \cdot h_{i_l}. \tag{2.1.5}$$


$n_l^{\mathbf{i}}$
$\mu_{w_{l-1}^{\mathbf{i}}}$
$\mu_{w_l^{\mathbf{i}}} = \mu_{w_{l-1}^{\mathbf{i}} s_{i_l}}$

Definition 2.1.3. A GGMS datum $\mu_\bullet = (\mu_w)_{w \in W}$ is said to be a Mirković-Vilonen (MV) datum if it satisfies the following condition:

(1) If two reduced words $\mathbf{i}, \mathbf{j} \in R(w_0)$ are related by a 2-move as in (2.1.3), then $n_l^{\mathbf{i}} = n_l^{\mathbf{j}}$ for all $1 \le l \le m$ with $l \neq k+1,\ k+2$, and $n_{k+1}^{\mathbf{i}} = n_{k+2}^{\mathbf{j}}$, $n_{k+2}^{\mathbf{i}} = n_{k+1}^{\mathbf{j}}$.


$\mu_{w_{k+2}^{\mathbf{i}}} = \mu_{w_k^{\mathbf{i}} s_i s_j} = \mu_{w_k^{\mathbf{j}} s_j s_i} = \mu_{w_{k+2}^{\mathbf{j}}}$
$n_{k+2}^{\mathbf{i}}$
$n_{k+2}^{\mathbf{j}}$
$\mu_{w_{k+1}^{\mathbf{i}}} = \mu_{w_k^{\mathbf{i}} s_i}$
$\mu_{w_{k+1}^{\mathbf{j}}} = \mu_{w_k^{\mathbf{j}} s_j}$
$n_{k+1}^{\mathbf{i}}$
$n_{k+1}^{\mathbf{j}}$
$\mu_{w_k^{\mathbf{i}}} = \mu_{w_k^{\mathbf{j}}}$

(2) If two reduced words $\mathbf{i}, \mathbf{j} \in R(w_0)$ are related by a 3-move as in (2.1.4), then $n_l^{\mathbf{i}} = n_l^{\mathbf{j}}$ for all $1 \le l \le m$ with $l \ne k+1,\ k+2,\ k+3$, and

$$\begin{cases} n_{k+1}^{\mathbf{j}} = n_{k+2}^{\mathbf{i}} + n_{k+3}^{\mathbf{i}} - \min\bigl(n_{k+1}^{\mathbf{i}},\, n_{k+3}^{\mathbf{i}}\bigr), \\ n_{k+2}^{\mathbf{j}} = \min\bigl(n_{k+1}^{\mathbf{i}},\, n_{k+3}^{\mathbf{i}}\bigr), \\ n_{k+3}^{\mathbf{j}} = n_{k+1}^{\mathbf{i}} + n_{k+2}^{\mathbf{i}} - \min\bigl(n_{k+1}^{\mathbf{i}},\, n_{k+3}^{\mathbf{i}}\bigr). \end{cases} \tag{2.1.6}$$


$\mu_{w^{\mathbf{i}}_{k+3}} = \mu_{w^{\mathbf{i}}_k s_i s_j s_i} = \mu_{w^{\mathbf{j}}_k s_j s_i s_j} = \mu_{w^{\mathbf{j}}_{k+3}}$
$n^{\mathbf{i}}_{k+3}$
$n^{\mathbf{j}}_{k+3}$
$\mu_{w^{\mathbf{i}}_{k+2}} = \mu_{w^{\mathbf{i}}_k s_i s_j}$
$\mu_{w^{\mathbf{j}}_{k+2}} = \mu_{w^{\mathbf{j}}_k s_j s_i}$
$n^{\mathbf{i}}_{k+2}$
$n^{\mathbf{j}}_{k+2}$
$\mu_{w^{\mathbf{i}}_{k+1}} = \mu_{w^{\mathbf{i}}_k s_i}$
$\mu_{w^{\mathbf{j}}_{k+1}} = \mu_{w^{\mathbf{j}}_k s_j}$
$n^{\mathbf{i}}_{k+1}$
$n^{\mathbf{j}}_{k+1}$
$\mu_{w^{\mathbf{i}}_k} = \mu_{w^{\mathbf{j}}_k}$

The pseudo-Weyl polytope $P(\mu_\bullet)$ with GGMS datum $\mu_\bullet = (\mu_w)_{w\in W}$ (see (2.1.2)) is called a Mirković-Vilonen (MV) polytope if the GGMS datum $\mu_\bullet = (\mu_w)_{w\in W}$ is an MV datum. We denote by $\mathcal{MV}$ the set of all MV polytopes.

Remark 2.1.4. In the definition above of MV polytopes, we first introduced the notion of MV datum (which was not introduced in [Kam1], [Kam2]), and then called the associated pseudo-Weyl polytope (by (2.1.2)) an MV polytope. This definition of MV polytopes is equivalent to the one in [Kam1], [Kam2], which is easily seen by an argument in the proof of [Kam1, Proposition 5.4] (see the comment following [Kam1, Theorem 7.1]).

2.2 Crystals $\mathcal{MV}(\infty)$ and $\mathcal{MV}(\lambda)$. For $j \in I$ and an MV datum $\mu_\bullet = (\mu_w)_{w\in W}$, we denote by $f_j\mu_\bullet$ (resp., $e_j\mu_\bullet$ if $\mu_e \ne \mu_{s_j}$) a unique MV datum $\mu'_\bullet = (\mu'_w)_{w\in W}$ such that $\mu'_e = \mu_e - h_j$ (resp., $\mu'_e = \mu_e + h_j$), and such that $\mu'_w = \mu_w$ for all $w \in W$ with $s_j w < w$ (see [Kam2, Theorem 3.5] and its proof); note that $\mu'_{w_0} = \mu_{w_0}$ and $\mu'_{s_j} = \mu_{s_j}$.

Now, let $\mathcal{MV}(\infty)$ denote the set of MV polytopes $P = P(\mu_\bullet)$ with GGMS (hence MV) datum $\mu_\bullet = (\mu_w)_{w\in W}$ for which $\mu_{w_0} = 0 \in \mathfrak{h}_\mathbb{R}$; note that, by the length formula, $\mu_w \in Q_-^\vee := \sum_{j\in I} \mathbb{Z}_{\le 0} h_j$ for all $w \in W$. Following [Kam2, §§3.3, 3.5, and 3.6], we endow $\mathcal{MV}(\infty)$ with a crystal structure for $U_q(\mathfrak{g}^\vee)$ (due to Lusztig [Lu2] and Berenstein-Zelevinsky [BeZ]) as follows. Let $P = P(\mu_\bullet) \in \mathcal{MV}(\infty)$ be an MV polytope with GGMS datum $\mu_\bullet = (\mu_w)_{w\in W}$. The weight $\mathrm{wt}(P)$ of P is, by definition, equal to the vertex $\mu_e \in Q_-^\vee$. For each $j \in I$, we define the (lowering) Kashiwara operator $f_j : \mathcal{MV}(\infty) \cup \{\mathbf{0}\} \to \mathcal{MV}(\infty) \cup \{\mathbf{0}\}$ and the (raising) Kashiwara operator $e_j : \mathcal{MV}(\infty) \cup \{\mathbf{0}\} \to \mathcal{MV}(\infty) \cup \{\mathbf{0}\}$ by:

$$
\begin{aligned}
e_j\mathbf{0} = f_j\mathbf{0} &:= \mathbf{0}, \\
f_jP = f_jP(\mu_\bullet) &:= P(f_j\mu_\bullet), \\
e_jP = e_jP(\mu_\bullet) &:= \begin{cases} P(e_j\mu_\bullet) & \text{if } \mu_e \neq \mu_{s_j}, \\ \mathbf{0} & \text{otherwise,} \end{cases}
\end{aligned}
$$

where $\mathbf{0}$ is an additional element, not contained in $\mathcal{MV}(\infty)$. We set $\varepsilon_j(P) := \max\{N \geq 0 \mid e_j^N P \neq \mathbf{0}\}$, and $\varphi_j(P) := \langle \alpha_j, \mathrm{wt}(P)\rangle + \varepsilon_j(P)$.

Theorem 2.2.1 ([Kam2, §3.3 and §3.6]). *The set $\mathcal{MV}(\infty)$, together with the maps* wt, *e_j, f_j ($j \in I$), and ε_j, φ_j ($j \in I$) above, is a crystal for $U_q(\mathfrak{g}^\vee)$. Moreover, there exists a unique isomorphism $\Psi : \mathcal{B}(\infty) \xrightarrow{\sim} \mathcal{MV}(\infty)$ of crystals for $U_q(\mathfrak{g}^\vee)$.*

Remark 2.2.2. Define a GGMS datum $\mu_\bullet = (\mu_w)_{w\in W}$ by: $\mu_w = 0 \in \mathfrak{h}_\mathbb{R}$ for all $w \in W$. It is obvious that $\mu_\bullet$ is an MV datum, and the MV polytope $P_0 := P(\mu_\bullet)$ is an element of $\mathcal{MV}(\infty)$ whose weight is $0 \in \mathfrak{h}_\mathbb{R}$. Therefore, under the isomorphism Ψ of Theorem 2.2.1, the element $u_\infty \in \mathcal{B}(\infty)$ corresponding to the identity element $1 \in U_q^-(\mathfrak{g}^\vee)$ is sent to the MV polytope $P_0 \in \mathcal{MV}(\infty)$.

Let $\lambda \in X_*(T) \subset \mathfrak{h}_\mathbb{R}$ be a dominant coweight. Let $\mathcal{MV}(\lambda)$ denote the set of MV polytopes $P = P(\mu_\bullet)$ with GGMS (hence MV) datum $\mu_\bullet = (\mu_w)_{w\in W}$ such that $\mu_{w_0} = \lambda$ and such that P is contained in the convex hull $\mathrm{Conv}(W\cdot\lambda)$ of the W-orbit $W \cdot \lambda \subset \mathfrak{h}_\mathbb{R}$; note that, by the length formula, $\mu_w \in \lambda + Q_-^\vee$ for all $w \in W$. Following [Kam2, §6.2], we endow $\mathcal{MV}(\lambda)$ with a crystal structure for $U_q(\mathfrak{g}^\vee)$ as follows. Let $P = P(\mu_\bullet) \in \mathcal{MV}(\lambda)$ be an MV polytope with GGMS datum $\mu_\bullet = (\mu_w)_{w\in W}$. The weight $\mathrm{wt}(P)$ of P is, by definition, equal to the vertex $\mu_e \in \lambda + Q_-^\vee$. For each $j \in I$, we define the lowering Kashiwara operator $f_j : \mathcal{MV}(\lambda) \cup \{\mathbf{0}\} \to \mathcal{MV}(\lambda) \cup \{\mathbf{0}\}$ and the raising Kashiwara operator $e_j : \mathcal{MV}(\lambda) \cup \{\mathbf{0}\} \to \mathcal{MV}(\lambda) \cup \{\mathbf{0}\}$ by:

$$
\begin{aligned}
e_j\mathbf{0} = f_j\mathbf{0} &:= \mathbf{0}, \\
f_jP = f_jP(\mu_\bullet) &:= \begin{cases} P(f_j\mu_\bullet) & \text{if } P(f_j\mu_\bullet) \subset \mathrm{Conv}(W\cdot\lambda), \\ \mathbf{0} & \text{otherwise,} \end{cases} \\
e_jP = e_jP(\mu_\bullet) &:= \begin{cases} P(e_j\mu_\bullet) & \text{if } \mu_e \neq \mu_{s_j}, \\ \mathbf{0} & \text{otherwise,} \end{cases}
\end{aligned}
$$

where $\mathbf{0}$ is an additional element, not contained in $\mathcal{MV}(\lambda)$. We set $\varepsilon_j(P) := \max\{N \geq 0 \mid e_j^N P \neq \mathbf{0}\}$, and $\varphi_j(P) := \max\{N \geq 0 \mid f_j^N P \neq \mathbf{0}\}$.

Theorem 2.2.3 ([Kam2, Theorem 6.4]). *The set $\mathcal{MV}(\lambda)$, together with the maps* wt, *e_j, f_j ($j \in I$), and ε_j, φ_j ($j \in I$) above, is a crystal for $U_q(\mathfrak{g}^\vee)$. Moreover, there exists a unique isomorphism $\Psi_\lambda : \mathcal{B}(\lambda) \xrightarrow{\sim} \mathcal{MV}(\lambda)$ of crystals for $U_q(\mathfrak{g}^\vee)$.*

Remark 2.2.4. (1) Define a GGMS datum $\mu_\bullet = (\mu_w)_{w \in W}$ by: $\mu_w = \lambda \in X_*(T) \subset \mathfrak{h}_\mathbb{R}$ for all $w \in W$. It is obvious that $\mu_\bullet$ is an MV datum, and the MV polytope $P_\lambda := P(\mu_\bullet)$ is an element of $\mathcal{MV}(\lambda)$ whose weight is $\lambda \in X_*(T)$. Therefore, under the isomorphism Ψ_λ of Theorem 2.2.3, the highest weight element $u_\lambda \in \mathcal{B}(\lambda)$ of weight λ is sent to the MV polytope $P_\lambda \in \mathcal{MV}(\lambda)$.

(2) We know from [An] that the polytope $P_{w_0 \cdot \lambda} := \mathrm{Conv}(W \cdot \lambda)$ is an element of $\mathcal{MV}(\lambda)$, and is the lowest weight element of weight $w_0 \cdot \lambda \in X_*(T)$; in fact, the GGMS (hence MV) datum of $P_{w_0 \cdot \lambda}$ is given by: $\mu_w = ww_0 \cdot \lambda \in \mathfrak{h}_\mathbb{R}$, $w \in W$ (see [Kam1] and also §4.1 below). Therefore, under the isomorphism Ψ_λ of Theorem 2.2.3, the lowest weight element $u_{w_0 \cdot \lambda} \in \mathcal{B}(\lambda)$ of weight $w_0 \cdot \lambda$ is sent to the MV polytope $P_{w_0 \cdot \lambda} \in \mathcal{MV}(\lambda)$.

3 MV polytopes lying in a Demazure crystal and an opposite Demazure crystal.

We fix (once and for all) an arbitrary dominant coweight $\lambda \in X_*(T) \subset \mathfrak{h}_\mathbb{R}$.

3.1 Demazure crystals. Let $x \in W$. The Demazure module $V_x(\lambda)$ is defined to be the $U_q^+(\mathfrak{g}^\vee)$-submodule of $V(\lambda)$ generated by the one-dimensional weight space $V(\lambda)_{x \cdot \lambda} \subset V(\lambda)$ of weight $x \cdot \lambda \in X_*(T) \subset \mathfrak{h}_\mathbb{R}$. Recall from [Kas2] that the Demazure crystal $\mathcal{B}_x(\lambda)$ is a subset of $\mathcal{B}(\lambda)$ such that

$$V(\lambda) \supset V_x(\lambda) = \bigoplus_{b \in \mathcal{B}_x(\lambda)} \mathbb{C}(q) G_\lambda(b), \tag{3.1.1}$$

where $G_\lambda(b)$, $b \in \mathcal{B}(\lambda)$, form the lower global basis of $V(\lambda)$.

Remark 3.1.1. If x, $y \in W$ satisfies $x \cdot \lambda = y \cdot \lambda$, then we have $V_x(\lambda) = V_y(\lambda)$ since $V(\lambda)_{x \cdot \lambda} = V(\lambda)_{y \cdot \lambda}$. Therefore, it follows from (3.1.1) that $\mathcal{B}_x(\lambda) = \mathcal{B}_y(\lambda)$.

We know from [Kas2, Proposition 3.2.3] that the Demazure crystals $\mathcal{B}_x(\lambda)$, $x \in W$, are characterized by the inductive relations:

$$\mathcal{B}_e(\lambda) = \{u_\lambda\}, \tag{3.1.2}$$

$$\mathcal{B}_x(\lambda) = \bigcup_{N \geq 0} f_j^N \mathcal{B}_{s_j x}(\lambda) \setminus \{\mathbf{0}\} \quad \text{for } x \in W \text{ and } j \in I \text{ with } s_j x < x. \tag{3.1.3}$$

In addition, we know from [Kas2, Proposition 3.2.5] that there exists a unique family $\mathcal{B}_x(\infty)$, $x \in W$, of subsets of $\mathcal{B}(\infty)$ satisfying the inductive relations:

$$\mathcal{B}_e(\infty) = \{u_\infty\}, \tag{3.1.4}$$

$$\mathcal{B}_x(\infty) = \bigcup_{N \geq 0} f_j^N \mathcal{B}_{s_j x}(\infty) \quad \text{for } x \in W \text{ and } j \in I \text{ with } s_j x < x; \tag{3.1.5}$$

for each $x \in W$, the subset $\mathcal{B}_x(\infty) \subset \mathcal{B}(\infty)$ is also called the Demazure crystal associated to x.

3.2 Condition for an MV polytope to lie in a Demazure crystal. Let us fix an arbitrary $x \in W$, and denote by p the length $\ell(xw_0)$ of $xw_0 \in W$. For each reduced word $\mathbf{i} = (i_1, i_2, \dots, i_m) \in R(w_0)$, with $m = \ell(w_0)$, we set

$$S(xw_0, \mathbf{i}) = \left\{ (a_1, a_2, \dots, a_p) \in [1, m]^p \;\middle|\; \begin{array}{l} 1 \le a_1 < a_2 < \cdots < a_p \le m \\ s_{i_{a_1}} s_{i_{a_2}} \cdots s_{i_{a_p}} = xw_0 \end{array} \right\},$$

where $[1, m] := \{a \in \mathbb{Z} \mid 1 \le a \le m\}$. Note that for each sequence $(a_1, a_2, \dots, a_p) \in S(xw_0, \mathbf{i})$, $xw_0 = s_{i_{a_1}} s_{i_{a_2}} \cdots s_{i_{a_p}}$ is a reduced expression since $p = \ell(xw_0)$. Let $\mathcal{MV}_x(\lambda)$ (resp., $\mathcal{MV}_x(\infty)$) denote the subset of $\mathcal{MV}(\lambda)$ (resp., $\mathcal{MV}(\infty)$) consisting of those elements $P(\mu_\bullet)$ with GGMS datum $\mu_\bullet = (\mu_w)_{w \in W}$ which satisfy the condition:

(Dem.) for some $\mathbf{i} \in R(w_0)$, there exists a sequence $\mathbf{a} = (a_1, a_2, \dots, a_p) \in S(xw_0, \mathbf{i})$ such that $n^{\mathbf{i}}_{a_q} = n^{\mathbf{i}}_{a_q}(\mu_\bullet) = 0$ for all $1 \le q \le p$ (for the definition of $n^{\mathbf{i}}_l \in \mathbb{Z}_{\ge 0}$, $1 \le l \le m$, see (2.1.5)).

The following is the first main result of this paper; its proof will be given in the next subsection.

Theorem 3.2.1. *Keep the notation above.*

(1) *Under the isomorphism* $\Psi_\lambda : \mathcal{B}(\lambda) \xrightarrow{\sim} \mathcal{MV}(\lambda)$ *of Theorem 2.2.3, the Demazure crystal* $\mathcal{B}_x(\lambda) \subset \mathcal{B}(\lambda)$ *associated to* $x \in W$ *is mapped to* $\mathcal{MV}_x(\lambda)$*, that is,*

$$\Psi_\lambda(\mathcal{B}_x(\lambda)) = \mathcal{MV}_x(\lambda).$$

(2) *Under the isomorphism* $\Psi : \mathcal{B}(\infty) \xrightarrow{\sim} \mathcal{MV}(\infty)$ *of Theorem 2.2.1, the Demazure crystal* $\mathcal{B}_x(\infty) \subset \mathcal{B}(\infty)$ *associated to* $x \in W$ *is mapped to* $\mathcal{MV}_x(\infty)$*, that is,*

$$\Psi(\mathcal{B}_x(\infty)) = \mathcal{MV}_x(\infty).$$

For each reduced word $\mathbf{i} = (i_1, i_2, \dots, i_m) \in R(w_0)$, we set

$$\widehat{S}(xw_0, \mathbf{i}) = \left\{ (a_1, a_2, \dots, a_l) \in [1, m]^l \;\middle|\; \begin{array}{l} 0 \le l \le m \\ 1 \le a_1 < a_2 < \cdots < a_l \le m \\ s_{i_{a_1}} s_{i_{a_2}} \cdots s_{i_{a_l}} = xw_0 \end{array} \right\};$$

note that we can replace the condition "$0 \le l \le m$" in the definition above with "$p \le l \le m$" since $\ell(xw_0) = p$. It is obvious that $S(xw_0, \mathbf{i}) \subset \widehat{S}(xw_0, \mathbf{i})$, since the set $S(xw_0, \mathbf{i})$ is identical to the subset of $\widehat{S}(xw_0, \mathbf{i})$ consisting of those elements $(a_1, a_2, \dots, a_l)$ for which $l = p$. Also, for each sequence $(a_1, a_2, \dots, a_l) \in \widehat{S}(xw_0, \mathbf{i})$, there exists a subsequence of it which is contained in $S(xw_0, \mathbf{i})$ (see, for example, [MoP, Chap. 5, §4, Corollary 2]). Using these facts, we obtain immediately the following corollary of Theorem 3.2.1.

Corollary 3.2.2. *The Demazure crystal* $\mathcal{MV}_x(\lambda)$ *(resp.,* $\mathcal{MV}_x(\infty)$*) is identical to the subset of* $\mathcal{MV}(\lambda)$ *(resp.,* $\mathcal{MV}(\infty)$*) consisting of those elements* $P(\mu_\bullet) \in \mathcal{MV}(\lambda)$ *(resp.,* $\in \mathcal{MV}(\infty)$*) with GGMS datum* $\mu_\bullet = (\mu_w)_{w \in W}$ *which satisfy the following condition: for some* $\mathbf{i} \in R(w_0)$*, there exists a sequence* $(a_1, a_2, \dots, a_l) \in \widehat{S}(xw_0, \mathbf{i})$ *such that* $n^{\mathbf{i}}_{a_q} = n^{\mathbf{i}}_{a_q}(\mu_\bullet) = 0$ *for all* $1 \le q \le l$*.*

3.3 Proof of Theorem 3.2.1. We keep the notation and assumptions of §3.2. The following proposition plays a key role in the proof of Theorem 3.2.1.

Proposition 3.3.1. *Let* $P(\mu_\bullet)$ *be an MV polytope in* $\mathcal{MV}_x(\lambda)$ *or* $\mathcal{MV}_x(\infty)$ *with GGMS datum* $\mu_\bullet = (\mu_w)_{w \in W}$*. Then, for every* $\mathbf{j} \in R(w_0)$*, there exists a sequence* $\mathbf{b} = (b_1, b_2, \dots, b_p) \in S(xw_0, \mathbf{j})$ *such that* $n^{\mathbf{j}}_{b_q} = n^{\mathbf{j}}_{b_q}(\mu_\bullet) = 0$ *for all* $1 \le q \le p$*.*

Proof. In view of Remark 2.1.2 along with the definitions of $\mathcal{MV}_x(\lambda)$ and $\mathcal{MV}_x(\infty)$, it suffices to show the following claim.

Claim. *Let* $\mathbf{i} \in R(w_0)$ *be such that there exists a sequence* $\mathbf{a} = (a_1, a_2, \dots, a_p) \in S(xw_0, \mathbf{i})$ *for which* $n^{\mathbf{i}}_{a_q} = n^{\mathbf{i}}_{a_q}(\mu_\bullet) = 0$ *for all* $1 \le q \le p$*. Assume that* $\mathbf{j} \in R(w_0)$ *is related to* $\mathbf{i} \in R(w_0)$ *by a 2-move or a 3-move. Then, there exists a sequence* $\mathbf{b} = (b_1, b_2, \dots, b_p) \in S(xw_0, \mathbf{j})$ *for which* $n^{\mathbf{j}}_{b_q} = 0$ *for all* $1 \le q \le p$*.*

Proof of Claim. We give a proof in the case that $\mathbf{i}$ and $\mathbf{j}$ are related by a 3-move as in (2.1.4), i.e.,

$$\begin{aligned} \mathbf{i} &= (i_1, \dots, i_k, i, j, i, i_{k+4}, \dots, i_m), \\ \mathbf{j} &= (i_1, \dots, i_k, j, i, j, i_{k+4}, \dots, i_m) \end{aligned}$$

for some indices $i, j \in I$ with $a_{ij} = a_{ji} = -1$ and an integer $0 \le k \le m-3$; the proof for the case of 2-move is similar (or, even simpler). Now, for $0 \le l_1, l_2 \le m$, we denote by $[l_1, l_2] \cap \mathbf{a}$ the subsequence of $\mathbf{a}$ consisting of those a_q's such that $a_q \in [l_1, l_2] := \{ l \in \mathbb{Z} \mid l_1 \le l \le l_2 \}$; by convention, we set $[l_1, l_2] \cap \mathbf{a} := \emptyset$ if there is no a_q such that $l_1 \le a_q \le l_2$. Then we have

$$\begin{aligned} &[1, k] \cap \mathbf{a} = (a_1, \dots, a_{u_0}), \\ &[k+1, k+3] \cap \mathbf{a} = (a_{u_0+1}, \dots, a_{u_1}), \\ &[k+4, m] \cap \mathbf{a} = (a_{u_1+1}, \dots, a_p) \end{aligned}$$

for $0 \le u_0 \le u_1 \le p$. We set $\mathbf{a}' := [k+1, k+3] \cap \mathbf{a}$. Since $xw_0 = s_{i_{a_1}} s_{i_{a_2}} \cdots s_{i_{a_p}}$ is a reduced expression and $i_{k+1} = i_{k+3} = i$, the subsequence $\mathbf{a}'$ cannot be equal to $(k+1, k+3)$. Namely, the subsequence $\mathbf{a}'$ is equal to one of the following: $\emptyset$, $(k+1)$, $(k+2)$, $(k+3)$, $(k+1, k+2)$, $(k+2, k+3)$, $(k+1, k+2, k+3)$. Note that if $\mathbf{a}' = (k+2)$ (and hence $n^{\mathbf{i}}_{k+2} = 0$), then $n^{\mathbf{j}}_{k+1} = 0$ or $n^{\mathbf{j}}_{k+3} = 0$ by (2.1.6). We define a strictly increasing sequence $\mathbf{b} = (b_1, b_2, \dots, b_p) \in [1, m]^p$ as follows:

$$b_q = a_q \quad \text{for } 1 \le q \le u_0 \text{ and } u_1 + 1 \le q \le p,$$

$$(b_{u_0+1},\, \dots,\, b_{u_1}) = \begin{cases} \emptyset & \text{if } \mathbf{a}' = \emptyset, \\ (k+2) & \text{if } \mathbf{a}' = (k+1) \text{ or } (k+3), \\ (k+1) & \text{if } \mathbf{a}' = (k+2) \text{ and } n^{\mathbf{j}}_{k+1} = 0, \\ (k+3) & \text{if } \mathbf{a}' = (k+2) \text{ and } n^{\mathbf{j}}_{k+1} \neq 0,\, n^{\mathbf{j}}_{k+3} = 0, \\ (k+2,\, k+3) & \text{if } \mathbf{a}' = (k+1,\, k+2), \\ (k+1,\, k+2) & \text{if } \mathbf{a}' = (k+2,\, k+3), \\ (k+1,\, k+2,\, k+3) & \text{if } \mathbf{a}' = (k+1,\, k+2,\, k+3). \end{cases}$$

Then, it follows that $\mathbf{b} \in S(xw_0,\, \mathbf{j})$. Also, by using (2.1.6), we can easily verify that $n^{\mathbf{j}}_{b_q} = 0$ for all $1 \le q \le p$. This proves the claim, completing the proof of the proposition. $\square$

Remark 3.3.2. By Proposition 3.3.1, we can replace the phrase "for some $\mathbf{i} \in R(w_0)$" in the condition (Dem.) (and in Corollary 3.2.2) for $\mathcal{MV}_x(\lambda)$ and $\mathcal{MV}_x(\infty)$ with the phrase "for every $\mathbf{i} \in R(w_0)$".

Proposition 3.3.3. (1) *We have* $\mathcal{MV}_e(\lambda) = \{P_\lambda\}$ *and* $\mathcal{MV}_e(\infty) = \{P_0\}$.

(2) *Let* $x \in W$ *and* $j \in I$ *be such that* $s_j x < x$. *Then, we have*

$$\mathcal{MV}_x(\lambda) = \bigcup_{N \ge 0} f_j^N \mathcal{MV}_{s_j x}(\lambda) \setminus \{\mathbf{0}\}, \tag{3.3.1}$$

$$\mathcal{MV}_x(\infty) = \bigcup_{N \ge 0} f_j^N \mathcal{MV}_{s_j x}(\infty). \tag{3.3.2}$$

Proof. (1) It is obvious from the definitions of $\mathcal{MV}_e(\lambda)$ and $\mathcal{MV}_e(\infty)$ that $P_\lambda \in \mathcal{MV}_e(\lambda)$ and $P_0 \in \mathcal{MV}_e(\infty)$. Now, let $P = P(\mu_\bullet) \in \mathcal{MV}_e(\lambda)$ (resp., $P = P(\mu_\bullet) \in \mathcal{MV}_e(\infty)$) be an MV polytope with GGMS datum $\mu_\bullet = (\mu_w)_{w \in W}$. Note that the set $S(ew_0,\, \mathbf{i}) = S(w_0,\, \mathbf{i})$ consists of the single element $(1,\, 2,\, \dots,\, m)$ for all $\mathbf{i} = (i_1,\, i_2,\, \dots,\, i_m) \in R(w_0)$. Therefore, we see from Proposition 3.3.1 that $n^{\mathbf{i}}_l(\mu_\bullet) = 0$ for all $\mathbf{i} \in R(w_0)$ and $1 \le l \le m$, which implies that $\mu_{w^{\mathbf{i}}_0} = \mu_{w^{\mathbf{i}}_1} = \cdots = \mu_{w^{\mathbf{i}}_m} = \mu_{w_0} = \lambda$ (resp., $= 0$) for all $\mathbf{i} \in R(w_0)$. Here we recall the well-known fact (see, for example, [BjB, Proposition 3.1.2]) that for each $w \in W$, there exist some $\mathbf{i} \in R(w_0)$ and an integer $0 \le l \le m$ such that $w = w^{\mathbf{i}}_l$. From this fact, it follows that $\mu_w = \lambda$ (resp., $\mu_w = 0$) for all $w \in W$, and hence $P = P_\lambda$ (resp., $P = P_0$). Thus we have shown part (1).

(2) We denote by p the length $\ell(xw_0)$ of xw_0 as in §3.2. Then the length $\ell(s_j xw_0)$ of $s_j xw_0$ is equal to $p+1$, since $s_j x < x$ by assumption. We take and fix $\mathbf{i} = (i_1,\, i_2,\, \dots,\, i_m) \in R(w_0)$ such that $i_1 = j$.

First we show the inclusion $\supset$ of (3.3.1) (resp., (3.3.2)). Let $P = P(\mu_\bullet) \in \mathcal{MV}_{s_j x}(\lambda)$ (resp., $P = P(\mu_\bullet) \in \mathcal{MV}_{s_j x}(\infty)$) be an MV polytope with GGMS datum $\mu_\bullet = (\mu_w)_{w \in W}$. Assume that $f_j^N P \neq \mathbf{0}$ for some $N \ge 0$, and let $\mu'_\bullet = (\mu'_w)_{w \in W}$ denote the GGMS (hence

MV) datum of the MV polytope $f_j^N P$, i.e., $f_j^N P = P(\mu'_\bullet)$. In order to show that $f_j^N P = P(\mu'_\bullet) \in \mathcal{MV}_x(\lambda)$ (resp., $\in \mathcal{MV}_x(\infty)$), we define a sequence $\mathbf{b} = (b_1, b_2, \dots, b_p) \in S(xw_0, \mathbf{i})$ as follows. By Proposition 3.3.1, there exists a sequence $\mathbf{a} = (a_1, a_2, \dots, a_{p+1}) \in S(s_j xw_0, \mathbf{i})$ such that $n^{\mathbf{i}}_{a_q}(\mu_\bullet) = 0$ for all $1 \le q \le p+1$; recall that $s_{i_{a_1}} s_{i_{a_2}} \cdots s_{i_{a_{p+1}}}$ is a reduced expression of $s_j xw_0$ by the definition of $S(s_j xw_0, \mathbf{i})$. Since $s_j xw_0 > xw_0$, it follows from the "exchange condition" (see, for example, [MoP, Chap. 5, §3, Proposition 2]) that xw_0 has a reduced expression of the form: $xw_0 = s_{i_{a_1}} \cdots s_{i_{a_{q-1}}} s_{i_{a_{q+1}}} \cdots s_{i_{a_{p+1}}}$ for some (uniquely determined) $1 \le q \le p+1$. Here we note that if $a_1 = 1$, then $q = 1$ (i.e., $s_{i_{a_1}}$ is removed) since $i_{a_1} = i_1 = j$ and $s_j xw_0 > xw_0$. Thus, the sequence $(a_1, \dots, a_{q-1}, a_{q+1}, \dots, a_{p+1})$ obtained from the sequence $\mathbf{a}$ by removing a_q is an element of $S(xw_0, \mathbf{i})$. We now define $\mathbf{b} = (b_1, b_2, \dots, b_p)$ to be this sequence $(a_1, \dots, a_{q-1}, a_{q+1}, \dots, a_{p+1}) \in S(xw_0, \mathbf{i})$. It follows that $b_q \ge 2$ for all $1 \le q \le p$.

Claim 1. *We have* $n^{\mathbf{i}}_{b_q}(\mu'_\bullet) = 0$ *for all* $1 \le q \le p$.

Proof of Claim 1. Note that $s_j w^{\mathbf{i}}_l < w^{\mathbf{i}}_l$ for all $1 \le l \le m$, since $\mathbf{i} = (i_1, i_2, \dots, i_m) \in R(w_0)$ is such that $i_1 = j$. Therefore, it follows from the definition of the Kashiwara operator f_j that $\mu'_{w^{\mathbf{i}}_l} = \mu_{w^{\mathbf{i}}_l}$ for all $1 \le l \le m$. Hence we have $n^{\mathbf{i}}_l(\mu'_\bullet) = n^{\mathbf{i}}_l(\mu_\bullet)$ for all $2 \le l \le m$. Because $b_q \ge 2$ for all $1 \le q \le p$, and $n^{\mathbf{i}}_{a_q}(\mu_\bullet) = 0$ for all $1 \le q \le p+1$, we deduce from the definition of the sequence $\mathbf{b}$ that $n^{\mathbf{i}}_{b_q}(\mu'_\bullet) = 0$ for all $1 \le q \le p$. This proves the claim. ▌

It follows from Claim 1 that $f_j^N P = P(\mu'_\bullet) \in \mathcal{MV}_x(\lambda)$ (resp., $\in \mathcal{MV}_x(\infty)$), and hence the inclusion $\supset$ of (3.3.1) (resp., (3.3.2)) is verified.

The inclusion $\subset$ of (3.3.1) (resp., (3.3.2)) can be shown similarly. Let $P = P(\mu_\bullet) \in \mathcal{MV}_x(\lambda)$ (resp., $P = P(\mu_\bullet) \in \mathcal{MV}_x(\infty)$) be an MV polytope with GGMS datum $\mu_\bullet = (\mu_w)_{w \in W}$. Let $\mu'_\bullet = (\mu'_w)_{w \in W}$ denote the GGMS (hence MV) datum of the MV polytope $e_j^{\max} P := e_j^{\varepsilon_j(P)} P$, i.e., $e_j^{\max} P = P(\mu'_\bullet)$. We define a sequence $\mathbf{b} = (b_1, b_2, \dots, b_{p+1}) \in S(s_j xw_0, \mathbf{i})$ as follows. By Proposition 3.3.1, there exists a sequence $\mathbf{a} = (a_1, a_2, \dots, a_p) \in S(xw_0, \mathbf{i})$ such that $n^{\mathbf{i}}_{a_q}(\mu_\bullet) = 0$ for all $1 \le q \le p$. Note that $a_1 \ge 2$, since $s_j xw_0 > xw_0$ and $\mathbf{i} = (i_1, i_2, \dots, i_m) \in R(w_0)$ is such that $i_1 = j$. We now define $\mathbf{b} = (b_1, b_2, \dots, b_{p+1})$ to be the sequence $\mathbf{b} = (1, a_1, a_2, \dots, a_p)$. Then it follows immediately that $\mathbf{b} \in S(s_j xw_0, \mathbf{i})$ since $i_1 = j$.

Claim 2. *We have* $n^{\mathbf{i}}_{b_q}(\mu'_\bullet) = 0$ *for all* $1 \le q \le p+1$.

Proof of Claim 2. Note that $e_j(e_j^{\max} P) = \mathbf{0}$ by the definition. Hence it follows from the definition of the Kashiwara operator e_j that $\mu'_{w^{\mathbf{i}}_0} = \mu'_e = \mu'_{s_j} = \mu'_{w^{\mathbf{i}}_1}$. Also, it follows from the definition of e_j that $\mu'_{w^{\mathbf{i}}_l} = \mu_{w^{\mathbf{i}}_l}$ for all $1 \le l \le m$, since $i_1 = j$ implies $s_j w^{\mathbf{i}}_l < w^{\mathbf{i}}_l$. Therefore, we have $n^{\mathbf{i}}_{b_1}(\mu'_\bullet) = n^{\mathbf{i}}_1(\mu'_\bullet) = 0$, and $n^{\mathbf{i}}_l(\mu'_\bullet) = n^{\mathbf{i}}_l(\mu_\bullet)$ for all $2 \le l \le m$. Because $a_q \ge a_1 \ge 2$ for all $1 \le q \le p$, and $n^{\mathbf{i}}_{a_q}(\mu_\bullet) = 0$ for all $1 \le q \le p$, we deduce that

$n^{\mathbf{i}}_{b_q}(\mu'_\bullet) = n^{\mathbf{i}}_{a_{q-1}}(\mu'_\bullet) = n^{\mathbf{i}}_{a_{q-1}}(\mu_\bullet) = 0$ for all $2 \le q \le p+1$. Hence we have $n^{\mathbf{i}}_{b_q}(\mu'_\bullet) = 0$ for all $1 \le q \le p+1$. This proves the claim. ▌

It follows from Claim 2 that $e_j^{\max} P = P(\mu'_\bullet) \in \mathcal{MV}_{s_jx}(\lambda)$, (resp., $\in \mathcal{MV}_{s_jx}(\infty)$). Therefore, we conclude that $P \in f_j^{\varepsilon_j(P)} \mathcal{MV}_{s_jx}(\lambda)$ (resp., $\in f_j^{\varepsilon_j(P)} \mathcal{MV}_{s_jx}(\infty)$), which implies the inclusion $\subset$ of (3.3.1) (resp., (3.3.2)). Thus, we have shown (3.3.1) and (3.3.2), thereby completing the proof of the proposition. □

Part (1) of Theorem 3.2.1 follows immediately by combining (3.1.2), (3.1.3) and the corresponding assertions for $\mathcal{MV}_x(\lambda)$, $x \in W$, in Proposition 3.3.3; also, recall Remark 2.2.4 (1). Similarly, part (2) of Theorem 3.2.1 follows immediately by combining (3.1.4), (3.1.5) and the corresponding assertions for $\mathcal{MV}_x(\infty)$, $x \in W$, in Proposition 3.3.3; also, recall Remark 2.2.2.

3.4 Opposite Demazure crystals. Let $x \in W$. The opposite Demazure module $V^x(\lambda)$ is defined to be the $U_q^-(\mathfrak{g}^\vee)$-submodule of $V(\lambda)$ generated by the one-dimensional weight space $V(\lambda)_{x\cdot\lambda} \subset V(\lambda)$ of weight $x \cdot \lambda \in X_*(T) \subset \mathfrak{h}_\mathbb{R}$. Recall from [Kas2] that the opposite Demazure crystal $\mathcal{B}^x(\lambda)$ is a subset of $\mathcal{B}(\lambda)$ such that

$$V(\lambda) \supset V^x(\lambda) = \bigoplus_{b \in \mathcal{B}^x(\lambda)} \mathbb{C}(q) G_\lambda(b), \tag{3.4.1}$$

where $G_\lambda(b)$, $b \in \mathcal{B}(\lambda)$, form the lower global basis of $V(\lambda)$.

Remark 3.4.1. If $x, y \in W$ satisfies $x \cdot \lambda = y \cdot \lambda$, then we have $V^x(\lambda) = V^y(\lambda)$ since $V(\lambda)_{x\cdot\lambda} = V(\lambda)_{y\cdot\lambda}$. Therefore, it follows from (3.4.1) that $\mathcal{B}^x(\lambda) = \mathcal{B}^y(\lambda)$.

We know from [Kas2, §4] that the opposite Demazure crystals $\mathcal{B}^x(\lambda)$, $x \in W$, are characterized by the (descending) inductive relations:

$$\mathcal{B}^{w_0}(\lambda) = \{u_{w_0\cdot\lambda}\}, \tag{3.4.2}$$

$$\mathcal{B}^x(\lambda) = \bigcup_{N \ge 0} e_j^N \mathcal{B}^{s_jx}(\lambda) \setminus \{\mathbf{0}\} \quad \text{for } x \in W \text{ and } j \in I \text{ with } s_jx > x. \tag{3.4.3}$$

Furthermore, we see from [Kas2, Proposition 4.2 (i)] and (3.4.3) that for $x \in W$ and $j \in I$ with $s_jx > x$, $b \in \mathcal{B}^x(\lambda)$ holds if and only if $f_j^{\max} b \in \mathcal{B}^{s_jx}(\lambda)$ holds, where for $b \in \mathcal{B}(\lambda)$ and $j \in I$, we set $f_j^{\max} b := f_j^{\varphi_j(b)} b$, with $\varphi_j(b) := \max\{N \ge 0 \mid f_j^N b \ne \mathbf{0}\}$. Using this fact successively, we obtain the following lemma (cf. [Kas4, Proposition 9.1.3 (2)] for a similar result for Demazure crystals).

Lemma 3.4.2. *Let $x \in W$, and let $i_1, i_2, \ldots, i_p$ be a sequence of elements in I such that $\ell(s_{i_p} \cdots s_{i_2} s_{i_1} x) = \ell(x) + p$ and $s_{i_p} \cdots s_{i_2} s_{i_1} x = w_0$. An element $b \in \mathcal{B}(\lambda)$ is contained in the opposite Demazure crystal $\mathcal{B}^x(\lambda)$ associated to x if and only if $f_{i_p}^{\max} \cdots f_{i_2}^{\max} f_{i_1}^{\max} b = u_{w_0\cdot\lambda}$.*

3.5 Condition for an MV polytope to lie in an opposite Demazure crystal. Let us fix an arbitrary $x \in W$, and denote by p the length $\ell(xw_0)$ of $xw_0 \in W$. Let $\mathcal{MV}^x(\lambda)$ denote the subset of $\mathcal{MV}(\lambda)$ consisting of those elements $P(\mu_\bullet) \in \mathcal{MV}(\lambda)$ with GGMS datum $\mu_\bullet = (\mu_w)_{w \in W}$ which satisfy the condition:

(Op. Dem.) for some $\mathbf{i} = (i_1, i_2, \dots, i_m) \in R(w_0)$ such that $w_p^{\mathbf{i}} = s_{i_1} s_{i_2} \cdots s_{i_p} = xw_0$, there holds $\mu_{w_l^{\mathbf{i}}} = w_l^{\mathbf{i}} w_0 \cdot \lambda$ for all $p \le l \le m$.

The following is the second main result of this paper.

Theorem 3.5.1. *Under the isomorphism $\Psi_\lambda : \mathcal{B}(\lambda) \xrightarrow{\sim} \mathcal{MV}(\lambda)$ of Theorem 2.2.3, the opposite Demazure crystal $\mathcal{B}^x(\lambda) \subset \mathcal{B}(\lambda)$ associated to $x \in W$ is mapped to $\mathcal{MV}^x(\lambda)$, that is,*

$$\Psi_\lambda(\mathcal{B}^x(\lambda)) = \mathcal{MV}^x(\lambda).$$

Proof. First, we prove that $\Psi_\lambda(\mathcal{B}^x(\lambda)) \subset \mathcal{MV}^x(\lambda)$. Let $P = P(\mu_\bullet) \in \Psi_\lambda(\mathcal{B}^x(\lambda))$ be an MV polytope with GGMS datum $\mu_\bullet = (\mu_w)_{w \in W}$, and take an arbitrary $\mathbf{i} = (i_1, i_2, \dots, i_m) \in R(w_0)$ such that $w_p^{\mathbf{i}} = s_{i_1} s_{i_2} \cdots s_{i_p} = xw_0$. We will show by descending induction on l that $\mu_{w_l^{\mathbf{i}}} = w_l^{\mathbf{i}} w_0 \cdot \lambda$ for every $p \le l \le m$. Since $\mu_{w_m^{\mathbf{i}}} = \mu_{w_0} = \lambda = w_0 w_0 \cdot \lambda = w_m^{\mathbf{i}} w_0 \cdot \lambda$, the assertion holds when $l = m$. For $p + 1 \le l \le m$, we have

$$\begin{aligned}
\mu_{w_{l-1}^{\mathbf{i}}} &= \mu_{w_l^{\mathbf{i}}} - n_l^{\mathbf{i}} w_{l-1}^{\mathbf{i}} \cdot h_{i_l} \quad \text{by the length formula} \\
&= w_l^{\mathbf{i}} w_0 \cdot \lambda - n_l^{\mathbf{i}} w_{l-1}^{\mathbf{i}} \cdot h_{i_l} \quad \text{by the induction hypothesis} \\
&= w_{l-1}^{\mathbf{i}} s_{i_l} w_0 \cdot \lambda - n_l^{\mathbf{i}} w_{l-1}^{\mathbf{i}} \cdot h_{i_l} \\
&= w_{l-1}^{\mathbf{i}} \cdot \Bigl(w_0 \cdot \lambda - \langle \alpha_{i_l}, w_0 \cdot \lambda \rangle h_{i_l} \Bigr) - n_l^{\mathbf{i}} w_{l-1}^{\mathbf{i}} \cdot h_{i_l} \\
&= w_{l-1}^{\mathbf{i}} w_0 \cdot \lambda - \bigl\{ \langle \alpha_{i_l}, w_0 \cdot \lambda \rangle + n_l^{\mathbf{i}} \bigr\} w_{l-1}^{\mathbf{i}} \cdot h_{i_l}.
\end{aligned}$$

Therefore, in order to show that $\mu_{w_{l-1}^{\mathbf{i}}} = w_{l-1}^{\mathbf{i}} w_0 \cdot \lambda$, it suffices to prove the following claim.

Claim 1. *We have $n_l^{\mathbf{i}} = -\langle \alpha_{i_l}, w_0 \cdot \lambda \rangle$.*

Proof of Claim 1. As in [Kam2, §6.3], we set

$$\begin{aligned}
&N_1 := \varphi_{i_1}(P), \quad N_2 := \varphi_{i_2}(f_{i_1}^{\max} P), \quad N_3 := \varphi_{i_3}(f_{i_2}^{\max} f_{i_1}^{\max} P), \quad \dots\dots \\
&\qquad \dots\dots, \quad N_m := \varphi_{i_m}(f_{i_{m-1}}^{\max} \cdots f_{i_2}^{\max} f_{i_1}^{\max} P),
\end{aligned} \tag{3.5.1}$$

where for $P \in \mathcal{MV}(\lambda)$ and $j \in I$, we set $f_j^{\max} P := f_j^{\varphi_j(P)} P \in \mathcal{MV}(\lambda)$. Then we have $N_{p+1} = N_{p+2} = \cdots = N_m = 0$. Indeed, it follows from the equality $s_{i_1} s_{i_2} \cdots s_{i_p} = xw_0$ that $s_{i_p} \cdots s_{i_2} s_{i_1} x = (xw_0)^{-1} x = w_0$, and hence

$$\ell(s_{i_p} \cdots s_{i_2} s_{i_1} x) = \ell(w_0) = m = (m - p) + p = \ell(x) + p,$$

since $\ell(x) = \ell(w_0) - \ell(xw_0) = m - p$. Therefore, from Lemma 3.4.2, we deduce that

$$f_{i_p}^{\max} \cdots f_{i_2}^{\max} f_{i_1}^{\max} P = P_{w_0 \cdot \lambda}.$$

Because $P_{w_0 \cdot \lambda} \in \mathcal{MV}(\lambda)$ is the lowest weight element of weight $w_0 \cdot \lambda$ (recall Remark 2.2.4 (2)), we have $\varphi_j(P_{w_0 \cdot \lambda}) = 0$ for all $j \in I$, and hence $N_{p+1} = N_{p+2} = \cdots = N_m = 0$, as desired. In particular, $N_l = 0$ since $p + 1 \le l \le m$. Also, we know from [Kam2, Theorem 6.6] (and the equation preceding it) that

$$N_l = \frac{1}{2} \langle w_{l-1}^{\mathbf{i}} \cdot \alpha_{i_l},\, \mu_{w_{l-1}^{\mathbf{i}}} + \mu_{w_l^{\mathbf{i}}} \rangle.$$

Consequently, we obtain

$$\langle w_{l-1}^{\mathbf{i}} \cdot \alpha_{i_l},\, \mu_{w_{l-1}^{\mathbf{i}}} \rangle = -\langle w_{l-1}^{\mathbf{i}} \cdot \alpha_{i_l},\, \mu_{w_l^{\mathbf{i}}} \rangle. \tag{3.5.2}$$

Here we note that by the length formula $\mu_{w_l^{\mathbf{i}}} - \mu_{w_{l-1}^{\mathbf{i}}} = n_l^{\mathbf{i}} w_{l-1}^{\mathbf{i}} \cdot h_{i_l}$, there holds

$$\langle w_{l-1}^{\mathbf{i}} \cdot \alpha_{i_l},\, \mu_{w_l^{\mathbf{i}}} - \mu_{w_{l-1}^{\mathbf{i}}} \rangle = n_l^{\mathbf{i}} \langle w_{l-1}^{\mathbf{i}} \cdot \alpha_{i_l},\, w_{l-1}^{\mathbf{i}} \cdot h_{i_l} \rangle. \tag{3.5.3}$$

The left-hand side of (3.5.3) is equal to:

$$\begin{aligned}
\langle w_{l-1}^{\mathbf{i}} \cdot \alpha_{i_l},\, \mu_{w_l^{\mathbf{i}}} - \mu_{w_{l-1}^{\mathbf{i}}} \rangle &= \langle w_{l-1}^{\mathbf{i}} \cdot \alpha_{i_l},\, \mu_{w_l^{\mathbf{i}}} \rangle - \langle w_{l-1}^{\mathbf{i}} \cdot \alpha_{i_l},\, \mu_{w_{l-1}^{\mathbf{i}}} \rangle \\
&= 2 \langle w_{l-1}^{\mathbf{i}} \cdot \alpha_{i_l},\, \mu_{w_l^{\mathbf{i}}} \rangle \quad \text{by (3.5.2)} \\
&= 2 \langle w_{l-1}^{\mathbf{i}} \cdot \alpha_{i_l},\, w_l^{\mathbf{i}} w_0 \cdot \lambda \rangle \quad \text{by the induction hypothesis} \\
&= 2 \langle w_{l-1}^{\mathbf{i}} \cdot \alpha_{i_l},\, w_{l-1}^{\mathbf{i}} s_{i_l} w_0 \cdot \lambda \rangle \\
&= 2 \langle \alpha_{i_l},\, s_{i_l} w_0 \cdot \lambda \rangle \\
&= -2 \langle \alpha_{i_l},\, w_0 \cdot \lambda \rangle.
\end{aligned}$$

The right-hand side of (3.5.3) is equal to:

$$n_l^{\mathbf{i}} \langle w_{l-1}^{\mathbf{i}} \cdot h_{i_l},\, w_{l-1}^{\mathbf{i}} \cdot \alpha_{i_l} \rangle = n_l^{\mathbf{i}} \langle h_{i_l},\, \alpha_{i_l} \rangle = 2 n_l^{\mathbf{i}}.$$

Substituting these equalities into (3.5.3), we conclude that $n_l^{\mathbf{i}} = -\langle \alpha_{i_l},\, w_0 \cdot \lambda \rangle$. This proves the claim. ∎

Thus, we have shown that $\mu_{w_l^{\mathbf{i}}} = w_l^{\mathbf{i}} w_0 \cdot \lambda$ for all $p \le l \le m$. This implies that $P \in \mathcal{MV}^x(\lambda)$, and hence that $\Psi_\lambda(\mathcal{B}^x(\lambda)) \subset \mathcal{MV}^x(\lambda)$.

Next, we prove the reverse inclusion $\Psi_\lambda(\mathcal{B}^x(\lambda)) \supset \mathcal{MV}^x(\lambda)$. Let $P = P(\mu_\bullet) \in \mathcal{MV}^x(\lambda)$ be an MV polytope with GGMS datum $\mu_\bullet = (\mu_w)_{w \in W}$. Since $P \in \mathcal{MV}^x(\lambda)$, there exists $\mathbf{i} = (i_1, i_2, \ldots, i_m) \in R(w_0)$ with $w_p^{\mathbf{i}} = s_{i_1} s_{i_2} \cdots s_{i_p} = x w_0$ such that $\mu_{w_l^{\mathbf{i}}} = w_l^{\mathbf{i}} w_0 \cdot \lambda$ for all $p \le l \le m$. Define nonnegative integers N_l, $1 \le l \le m$, as in (3.5.1):

$$N_l := \varphi_{i_l}(f_{i_{l-1}}^{\max} \cdots f_{i_2}^{\max} f_{i_1}^{\max} P).$$

Claim 2. *We have* $N_l = 0$ *for all* $p+1 \le l \le m$.

Proof of Claim 2. Let $p+1 \le l \le m$. We know from [Kam2, Theorem 6.6] (and the equation preceding it) that

$$N_l = \frac{1}{2}\langle w^{\mathbf{i}}_{l-1} \cdot \alpha_{i_l},\, \mu_{w^{\mathbf{i}}_{l-1}} + \mu_{w^{\mathbf{i}}_l}\rangle.$$

Using the equalities $\mu_{w^{\mathbf{i}}_{l-1}} = w^{\mathbf{i}}_{l-1}w_0 \cdot \lambda$ and $\mu_{w^{\mathbf{i}}_l} = w^{\mathbf{i}}_l w_0 \cdot \lambda$, we calculate as follows:

$$\begin{aligned}\langle w^{\mathbf{i}}_{l-1} \cdot \alpha_{i_l},\, \mu_{w^{\mathbf{i}}_{l-1}} + \mu_{w^{\mathbf{i}}_l}\rangle &= \langle w^{\mathbf{i}}_{l-1} \cdot \alpha_{i_l},\, w^{\mathbf{i}}_{l-1}w_0 \cdot \lambda\rangle + \langle w^{\mathbf{i}}_{l-1} \cdot \alpha_{i_l},\, w^{\mathbf{i}}_l w_0 \cdot \lambda\rangle \\ &= \langle w^{\mathbf{i}}_{l-1} \cdot \alpha_{i_l},\, w^{\mathbf{i}}_{l-1}w_0 \cdot \lambda\rangle + \langle w^{\mathbf{i}}_{l-1} \cdot \alpha_{i_l},\, w^{\mathbf{i}}_{l-1}s_{i_l} w_0 \cdot \lambda\rangle \\ &= \langle \alpha_{i_l},\, w_0 \cdot \lambda\rangle + \langle \alpha_{i_l},\, s_{i_l}w_0 \cdot \lambda\rangle \\ &= \langle \alpha_{i_l},\, w_0 \cdot \lambda\rangle - \langle \alpha_{i_l},\, w_0 \cdot \lambda\rangle = 0.\end{aligned}$$

Therefore, we obtain $N_l = 0$, as desired. ▮

We know (see, for example, [Kam2, Proposition 6.5]) that

$$f^{\max}_{i_m} \cdots f^{\max}_{i_2} f^{\max}_{i_1} P = f^{N_m}_{i_m} \cdots f^{N_2}_{i_2} f^{N_1}_{i_1} P = P_{w_0 \cdot \lambda}.$$

Here, by Claim 2, we have

$$f^{N_m}_{i_m} \cdots f^{N_2}_{i_2} f^{N_1}_{i_1} P = f^{N_p}_{i_p} \cdots f^{N_2}_{i_2} f^{N_1}_{i_1} P = f^{\max}_{i_p} \cdots f^{\max}_{i_2} f^{\max}_{i_1} P.$$

Hence we obtain $f^{\max}_{i_p} \cdots f^{\max}_{i_2} f^{\max}_{i_1} P = P_{w_0 \cdot \lambda}$. Since $s_{i_1}s_{i_2}\cdots s_{i_p} = xw_0$, we deduce that $s_{i_p}\cdots s_{i_2}s_{i_1}x = w_0$, and hence $\ell(s_{i_p}\cdots s_{i_2}s_{i_1}x) = \ell(x) + p$, as in the proof of Claim 1. Therefore, from Lemma 3.4.2, we conclude that $P \in \Psi_\lambda(\mathcal{B}^x(\lambda))$. Thus, we have shown that $\Psi_\lambda(\mathcal{B}^x(\lambda)) \supset \mathcal{MV}^x(\lambda)$, and hence that $\Psi_\lambda(\mathcal{B}^x(\lambda)) = \mathcal{MV}^x(\lambda)$. This completes the proof of the theorem. □

Remark 3.5.2. We see from the proof of Theorem 3.5.1 that we can replace the phrase "for some $\mathbf{i} = (i_1, i_2, \ldots, i_m) \in R(w_0)$" in the condition (Op. Dem.) for $\mathcal{MV}^x(\lambda)$ with the phrase "for every $\mathbf{i} = (i_1, i_2, \ldots, i_m) \in R(w_0)$".

4 Extremal MV polytopes and its relation with opposite Demazure crystals.

We fix (once and for all) an arbitrary dominant coweight $\lambda \in X_*(T) \subset \mathfrak{h}_\mathbb{R}$. For each $x \in W$, we denote by $P_{x\cdot\lambda}$ the image of the extremal element $u_{x\cdot\lambda} \in \mathcal{B}(\lambda)$ of weight $x\cdot\lambda \in X_*(T) \subset \mathfrak{h}_\mathbb{R}$ under the isomorphism $\Psi_\lambda : \mathcal{B}(\lambda) \xrightarrow{\sim} \mathcal{MV}(\lambda)$ of Theorem 2.2.3; we call $P_{x\cdot\lambda} \in \mathcal{MV}(\lambda)$ the extremal MV polytope of weight $x \cdot \lambda$. Recall (from [N, Lemma 4.2], for example) that for every reduced expression $x = s_{i_1}s_{i_2}\cdots s_{i_l}$ of x, we have $u_{x\cdot\lambda} = f^{\max}_{i_1} f^{\max}_{i_2} \cdots f^{\max}_{i_l} u_\lambda$. The aim of this section is to give an explicit description as a pseudo-Weyl polytope for the extremal MV polytopes $P_{x\cdot\lambda}$, $x \in W$, and to give a polytopal condition for an MV polytope $P \in \mathcal{MV}(\lambda)$ to lie in the opposite Demazure crystal $\mathcal{MV}^x(\lambda)$ for $x \in W$.

4.1 Explicit description of extremal MV polytopes. We fix $x \in W$, and denote by p the length $\ell(xw_0)$ of xw_0. Let $\mathbf{i} = (i_1, i_2, \dots, i_m)$ be an arbitrary element of $R(w_0)$, with $m = \ell(w_0)$. For a sequence $\mathbf{a} = (a_1, a_2, \dots, a_p) \in S(xw_0, \mathbf{i})$ and $0 \le l_1, l_2 \le m$, let $[l_1, l_2] \cap \mathbf{a}$ denote the subsequence of $\mathbf{a}$ consisting of those a_q's such that $a_q \in [l_1, l_2]$. Also, recall that the lexicographic ordering $\succeq$ on $S(xw_0, \mathbf{i})$ is defined as follows: $(a_1, a_2, \dots, a_p) \succ (b_1, b_2, \dots, b_p)$ if there exists some integer $1 \le q_0 \le p$ such that $a_q = b_q$ for all $1 \le q \le q_0 - 1$ and $a_{q_0} > b_{q_0}$; we denote by $\min S(xw_0, \mathbf{i})$ the minimum element of $S(xw_0, \mathbf{i})$ with respect to the lexicographic ordering.

We define a sequence $\xi_0^{\mathbf{i}}, \xi_1^{\mathbf{i}}, \dots, \xi_m^{\mathbf{i}}$ of elements of the W-orbit $W \cdot \lambda$ $(\subset \mathfrak{h}_{\mathbb{R}})$ inductively by the following formula:

$$\xi_m^{\mathbf{i}} = \lambda, \qquad \xi_{l-1}^{\mathbf{i}} = \begin{cases} \xi_l^{\mathbf{i}} & \text{if } l \text{ appears in } \min S(xw_0, \mathbf{i}), \\ s_{\beta_l^{\mathbf{i}}} \cdot \xi_l^{\mathbf{i}} & \text{otherwise} \end{cases} \tag{4.1.1}$$

for $1 \le l \le m$, where we set $\beta_l^{\mathbf{i}} := w_{l-1}^{\mathbf{i}} \cdot \alpha_{i_l}$ for $1 \le l \le m$, and denote by $s_\beta \in W$ the reflection with respect to a root β.

Remark 4.1.1. It is well-known (see, for example, [MoP, Chap. 5, §2, Lemma 2 and Proposition 3]) that $\beta_l^{\mathbf{i}}$, $1 \le l \le m$, exhaust all the positive roots.

Example 4.1.2. We know (see, for example, [BjB, Proposition 3.1.2]) that there exists $\mathbf{i} \in R(w_0)$ such that $w_p^{\mathbf{i}} = xw_0$. It is obvious that for this $\mathbf{i} \in R(w_0)$, $\min S(xw_0, \mathbf{i}) = (1, 2, \dots, p)$. Hence it follows from the definition that $\xi_0^{\mathbf{i}} = \xi_1^{\mathbf{i}} = \cdots = \xi_p^{\mathbf{i}}$ and $\xi_l^{\mathbf{i}} = s_{\beta_{l+1}^{\mathbf{i}}} s_{\beta_{l+2}^{\mathbf{i}}} \cdots s_{\beta_m^{\mathbf{i}}} \cdot \lambda$ for all $p \le l \le m$. Moreover, we see from (4.2.1) below that $\xi_l^{\mathbf{i}} = w_l^{\mathbf{i}} w_0 \cdot \lambda$ for all $p \le l \le m$.

Lemma 4.1.3. *We have* $\xi_l^{\mathbf{i}} - \xi_{l-1}^{\mathbf{i}} \in \mathbb{Z}_{\ge 0}(w_{l-1}^{\mathbf{i}} \cdot h_{i_l})$ *for every* $1 \le l \le m$.

The proof of this lemma will be given in §4.2. From this lemma, using [Kam1, Theorem 7.1], we see that there exists a unique MV datum $\mu_\bullet^{x,\mathbf{i}} = (\mu_w^{x,\mathbf{i}})_{w \in W}$ such that $\mu_{w_l^{\mathbf{i}}}^{x,\mathbf{i}} = \xi_l^{\mathbf{i}}$ for every $0 \le l \le m$.

Proposition 4.1.4. *Let* $\mathbf{j} \in R(w_0)$ *be another reduced word for* w_0*, and define an MV datum* $\mu_\bullet^{x,\mathbf{j}}$ *in the same way as above, with* $\mathbf{i}$ *replaced by* $\mathbf{j}$*. Then, we have* $\mu_\bullet^{x,\mathbf{i}} = \mu_\bullet^{x,\mathbf{j}}$.

The proof of this proposition will be given in §4.3. It follows from this proposition that the MV datum $\mu_\bullet^x = (\mu_w^x)_{w \in W} := \mu_\bullet^{x,\mathbf{i}}$ does not depend on the choice of $\mathbf{i} \in R(w_0)$. Moreover, the MV polytope $P(\mu_\bullet^x)$ associated to the MV datum $\mu_\bullet^x$ (by (2.1.2)) is an element of $\mathcal{MV}(\lambda)$. Indeed, recall that each $w \in W$ can be written as $w = w_l^{\mathbf{i}}$ for some $\mathbf{i} \in R(w_0)$ and an integer $0 \le l \le m$. Then it follows that $\mu_w^x = \mu_{w_l^{\mathbf{i}}}^x = \mu_{w_l^{\mathbf{i}}}^{x,\mathbf{i}} = \xi_l^{\mathbf{i}} \in W \cdot \lambda$ by the definition of $\mu_\bullet^x$. Since $w_0 = w_m^{\mathbf{i}}$ for all $\mathbf{i} \in R(w_0)$, and since $\xi_m^{\mathbf{i}} = \lambda$, we get $\mu_{w_0}^x = \lambda$. Also, because $\xi_l^{\mathbf{i}} \in W \cdot \lambda$ for all $\mathbf{i} \in R(w_0)$ and $0 \le l \le m$, we have

$$\begin{aligned} P(\mu_\bullet^x) &= \mathrm{Conv}\left\{\mu_w^x \mid w \in W\right\} \\ &= \mathrm{Conv}\left\{\xi_l^{\mathbf{i}} \mid \mathbf{i} \in R(w_0),\, 0 \le l \le m\right\} \subset \mathrm{Conv}(W \cdot \lambda). \end{aligned} \tag{4.1.2}$$

Theorem 4.1.5. (1) *The weight of the MV polytope* $P(\mu_\bullet^x) \in \mathcal{MV}(\lambda)$ *is equal to* $x \cdot \lambda$. *Therefore,* $P(\mu_\bullet^x)$ *is the extremal MV polytope* $P_{x\cdot\lambda}$ *of weight* $x \cdot \lambda$.

(2) *The extremal MV polytope* $P_{x\cdot\lambda} = P(\mu_\bullet^x)$ *of weight* $x \cdot \lambda$ *is identical to the convex hull* $\mathrm{Conv}(W_{\le x} \cdot \lambda)$ *in* $\mathfrak{h}_\mathbb{R}$ *of the set* $W_{\le x} \cdot \lambda$, *where* $W_{\le x}$ *denotes the subset* $\left\{z \in W \mid z \le x\right\}$ *of* W.

The proof of this theorem will be given in §4.4. By combining Theorems 3.5.1 and 4.1.5, we obtain a polytopal condition for an MV polytope $P \in \mathcal{MV}(\lambda)$ to lie in the opposite Demazure crystal $\mathcal{MV}^x(\lambda)$ for $x \in W$.

Remark 4.1.6. The statement of Theorem 4.1.5 (2) seems to be known to some experts although we were unable to find it explicitly stated in the literature (cf. [AP], and also [At, §4]). In fact, as was reported to us by Kato [Kat], the equality $P_{x\cdot\lambda} = \mathrm{Conv}(W_{\le x} \cdot \lambda)$ follows from the first equality in [MiV2, (3.6)] (more precisely, from [Ha, Lemma 3.2] or [NP, Lemme 5.2]) by using the geometry of the affine Grassmannian $\mathcal{G}r$ and the finer Bruhat decomposition:

$$G = \bigsqcup_{z \in W_{\min}^\lambda} N \dot{z} P_\lambda,$$

where P_λ ($\supset B$) is a standard parabolic subgroup of G determined by the set of roots α for which $\langle \alpha, \lambda \rangle \le 0$.

However, the equality $P_{x\cdot\lambda} = \mathrm{Conv}(W_{\le x} \cdot \lambda)$ is not enough for our purposes. To get an understanding of the extremal MV polytope $P_{x\cdot\lambda}$ as a pseudo-Weyl polytope, we need to identify explicitly the vertex μ_w of $P_{x\cdot\lambda}$ for each $w \in W$; such data can be obtained from our construction above of $P_{x\cdot\lambda}$. In particular, it follows immediately from Theorems 3.2.1 and 3.5.1 that $P_{x\cdot\lambda} \in \mathcal{MV}_x(\lambda) \cap \mathcal{MV}^x(\lambda)$.

4.2 Proof of Lemma 4.1.3. Keep the notation and assumptions of §4.1. Let $\mathbf{i} = (i_1, i_2, \ldots, i_m)$ be an arbitrary element of $R(w_0)$. For each $0 \le l \le m$, we denote by $[l+1, m] \setminus \min S(xw_0, \mathbf{i})$ the sequence obtained by enumerating (in increasing order) the integers in $[l+1, m]$ that do not appear in the sequence $\min S(xw_0, \mathbf{i})$. Now, we write the sequence $[l+1, m] \setminus \min S(xw_0, \mathbf{i})$ as $(b_1, b_2, \ldots, b_{t_l})$, and set

$$y_l^{\mathbf{i}} := s_{\beta_{b_1}^{\mathbf{i}}} s_{\beta_{b_2}^{\mathbf{i}}} \cdots s_{\beta_{b_{t_l}}^{\mathbf{i}}};$$

if all the integers in $[l+1, m]$ appear in the sequence $\min S(xw_0, \mathbf{i})$, then we set $[l+1, m] \setminus \min S(xw_0, \mathbf{i}) = \emptyset$, and $y_l^{\mathbf{i}} := e \in W$. Note that $\xi_l^{\mathbf{i}} = y_l^{\mathbf{i}} \cdot \lambda$ for all $0 \le l \le m$. Also, for each $0 \le l \le m$, we write the sequence $[l+1,\, m] \cap \min S(xw_0,\, \mathbf{i})$ as $(c_1,\, c_2,\, \ldots,\, c_{u_l})$, with $u_l + t_l = m - l$, and set

$$v_l^{\mathbf{i}} := s_{i_{c_1}} s_{i_{c_2}} \cdots s_{i_{c_{u_l}}};$$

if $[l+1,\, m] \cap \min S(xw_0,\, \mathbf{i}) = \emptyset$, then we set $v_l^{\mathbf{i}} := e \in W$.

Lemma 4.2.1. *With the notation above, we have* $y_l^{\mathbf{i}} = w_l^{\mathbf{i}} v_l^{\mathbf{i}} w_0^{-1}$ *for every* $0 \le l \le m$. *Hence, for every* $0 \le l \le m$, *we have*

$$\xi_l^{\mathbf{i}} = w_l^{\mathbf{i}} v_l^{\mathbf{i}} w_0^{-1} \cdot \lambda. \tag{4.2.1}$$

Proof. We show the lemma by descending induction on l. Assume first that $l = m$. Then we have $y_m^{\mathbf{i}} = e$ and $v_m^{\mathbf{i}} = e$ by definition. Also, from the definition, we have $w_m^{\mathbf{i}} = w_0$. Hence we get $w_m^{\mathbf{i}} v_m^{\mathbf{i}} w_0^{-1} = w_0 e w_0^{-1} = e = y_m^{\mathbf{i}}$. Assume now that $l < m$. If $l+1$ appears in the sequence $\min S(xw_0,\, \mathbf{i})$, then we have

$$\begin{aligned} &[l+1,\, m] \setminus \min S(xw_0,\, \mathbf{i}) = [l+2,\, m] \setminus \min S(xw_0,\, \mathbf{i}), \\ &[l+1,\, m] \cap \min S(xw_0,\, \mathbf{i}) = \Big(l+1,\, [l+2,\, m] \cap \min S(xw_0,\, \mathbf{i})\Big), \end{aligned}$$

and hence $y_l^{\mathbf{i}} = y_{l+1}^{\mathbf{i}}$ and $v_l^{\mathbf{i}} = s_{i_{l+1}} v_{l+1}^{\mathbf{i}}$. Therefore, we conclude that

$$\begin{aligned} y_l^{\mathbf{i}} = y_{l+1}^{\mathbf{i}} &= w_{l+1}^{\mathbf{i}} v_{l+1}^{\mathbf{i}} w_0^{-1} \quad \text{by the induction hypothesis} \\ &= w_l^{\mathbf{i}} s_{i_{l+1}} v_{l+1}^{\mathbf{i}} w_0^{-1} = w_l^{\mathbf{i}} s_{i_{l+1}} s_{i_{l+1}} v_l^{\mathbf{i}} w_0^{-1} = w_l^{\mathbf{i}} v_l^{\mathbf{i}} w_0^{-1}. \end{aligned}$$

If $l+1$ does not appear in the sequence $\min S(xw_0,\, \mathbf{i})$, then we have

$$\begin{aligned} &[l+1,\, m] \setminus \min S(xw_0,\, \mathbf{i}) = \Big(l+1,\, [l+2,\, m] \setminus \min S(xw_0,\, \mathbf{i})\Big), \\ &[l+1,\, m] \cap \min S(xw_0,\, \mathbf{i}) = [l+2,\, m] \cap \min S(xw_0,\, \mathbf{i}), \end{aligned}$$

and hence $y_l^{\mathbf{i}} = s_{\beta_{l+1}^{\mathbf{i}}} y_{l+1}^{\mathbf{i}}$ and $v_l^{\mathbf{i}} = v_{l+1}^{\mathbf{i}}$. By the induction hypothesis, we get

$$y_l^{\mathbf{i}} = s_{\beta_{l+1}^{\mathbf{i}}} y_{l+1}^{\mathbf{i}} = s_{\beta_{l+1}^{\mathbf{i}}} w_{l+1}^{\mathbf{i}} v_{l+1}^{\mathbf{i}} w_0^{-1}.$$

Since $s_{\beta_{l+1}^{\mathbf{i}}} w_{l+1}^{\mathbf{i}} = \big(w_l^{\mathbf{i}} s_{i_{l+1}} (w_l^{\mathbf{i}})^{-1}\big) w_{l+1}^{\mathbf{i}} = w_l^{\mathbf{i}}$ (note that $s_{i_{l+1}} (w_l^{\mathbf{i}})^{-1} w_{l+1}^{\mathbf{i}} = (w_l^{\mathbf{i}} s_{i_{l+1}})^{-1} w_{l+1}^{\mathbf{i}} = (w_{l+1}^{\mathbf{i}})^{-1} w_{l+1}^{\mathbf{i}} = e$), it follows that

$$y_l^{\mathbf{i}} = s_{\beta_{l+1}^{\mathbf{i}}} w_{l+1}^{\mathbf{i}} v_{l+1}^{\mathbf{i}} w_0^{-1} = w_l^{\mathbf{i}} v_{l+1}^{\mathbf{i}} w_0^{-1} = w_l^{\mathbf{i}} v_l^{\mathbf{i}} w_0^{-1}.$$

Thus we have shown that $y_l^{\mathbf{i}} = w_l^{\mathbf{i}} v_l^{\mathbf{i}} w_0^{-1}$ for all $0 \le l \le m$. This proves the lemma. □

Proof of Lemma 4.1.3. Let $\mathbf{i} = (i_1,\, i_2,\, \ldots,\, i_m)$ be an arbitrary element of $R(w_0)$, and let $1 \le l \le m$. First we observe that $w_{l-1}^{\mathbf{i}} \cdot h_{i_l} = (w_{l-1}^{\mathbf{i}} \cdot \alpha_{i_l})^{\vee} = (\beta_l^{\mathbf{i}})^{\vee}$, where $\beta^{\vee}$ denotes the dual root of a positive root β, and hence that

$$s_{\beta_l^{\mathbf{i}}} \cdot \xi_l^{\mathbf{i}} = \xi_l^{\mathbf{i}} - \langle \beta_l^{\mathbf{i}},\, \xi_l^{\mathbf{i}} \rangle (\beta_l^{\mathbf{i}})^{\vee} = \xi_l^{\mathbf{i}} - \langle \beta_l^{\mathbf{i}},\, \xi_l^{\mathbf{i}} \rangle (w_{l-1}^{\mathbf{i}} \cdot h_{i_l}).$$

Therefore, we see from the definition (4.1.1) that

$$\xi_l^{\mathbf{i}} - \xi_{l-1}^{\mathbf{i}} = \begin{cases} 0 & \text{if } l \text{ appears in } \min S(xw_0,\, \mathbf{i}), \\ \langle \beta_l^{\mathbf{i}},\, \xi_l^{\mathbf{i}} \rangle (w_{l-1}^{\mathbf{i}} \cdot h_{i_l}) & \text{otherwise.} \end{cases} \tag{4.2.2}$$

So, it suffices to show that $\langle \beta_l^{\mathbf{i}},\, \xi_l^{\mathbf{i}} \rangle \geq 0$ under the assumption that l does not appear in the sequence $\min S(xw_0,\, \mathbf{i})$. By (4.2.1), we have

$$\xi_l^{\mathbf{i}} = w_l^{\mathbf{i}} v_l^{\mathbf{i}} w_0^{-1} \cdot \lambda = w_l^{\mathbf{i}} \, (s_{i_{c_1}} s_{i_{c_2}} \cdots s_{i_{c_{u_l}}}) \, w_0^{-1} \cdot \lambda,$$

where, as above, $[l+1,\, m] \cap \min S(xw_0,\, \mathbf{i}) = (c_1,\, c_2,\, \ldots,\, c_{u_l})$. Hence we calculate:

$$\begin{aligned} \langle \beta_l^{\mathbf{i}},\, \xi_l^{\mathbf{i}} \rangle &= \big\langle w_{l-1}^{\mathbf{i}} \cdot \alpha_{i_l},\, w_l^{\mathbf{i}} \, (s_{i_{c_1}} s_{i_{c_2}} \cdots s_{i_{c_{u_l}}}) \, w_0^{-1} \cdot \lambda \big\rangle \\ &= \big\langle (w_l^{\mathbf{i}})^{-1} w_{l-1}^{\mathbf{i}} \cdot \alpha_{i_l},\, (s_{i_{c_1}} s_{i_{c_2}} \cdots s_{i_{c_{u_l}}}) \, w_0^{-1} \cdot \lambda \big\rangle \\ &= \big\langle s_{i_l} \cdot \alpha_{i_l},\, (s_{i_{c_1}} s_{i_{c_2}} \cdots s_{i_{c_{u_l}}}) \, w_0^{-1} \cdot \lambda \big\rangle \\ &= -\big\langle \alpha_{i_l},\, (s_{i_{c_1}} s_{i_{c_2}} \cdots s_{i_{c_{u_l}}}) \, w_0^{-1} \cdot \lambda \big\rangle \\ &= -\big\langle w_0 (s_{i_{c_{u_l}}} \cdots s_{i_{c_2}} s_{i_{c_1}}) \cdot \alpha_{i_l},\, \lambda \big\rangle. \end{aligned}$$

Since $\lambda \in X_*(T) \subset \mathfrak{h}_{\mathbb{R}}$ is a dominant coweight, it suffices to show that $(s_{i_{c_{u_l}}} \cdots s_{i_{c_2}} s_{i_{c_1}}) \cdot \alpha_{i_l}$ is a positive root. Suppose, contrary to our claim, that it is a negative root. Then, by the exchange condition, $s_{i_{c_1}} s_{i_{c_2}} \cdots s_{i_{c_{u_l}}}$ has a reduced expression of the form:

$$s_{i_l} s_{i_{c_1}} \cdots s_{i_{c_{r-1}}} s_{i_{c_{r+1}}} \cdots s_{i_{c_{u_l}}}$$

for some $1 \leq r \leq u_l$. We now write the sequence $[1,\, l] \cap \min S(xw_0,\, \mathbf{i})$ as:

$$[1,\, l] \cap \min S(xw_0,\, \mathbf{i}) = \big\{ d_1,\, d_2,\, \ldots,\, d_{p-u_l} \big\};$$

note that $d_{p-u_l} < l$ since l does not appear in $\min S(xw_0,\, \mathbf{i})$. Then, since $\ell(xw_0) = p = (p - u_l) + u_l$,

$$\begin{aligned} xw_0 &= s_{i_{d_1}} s_{i_{d_2}} \cdots s_{i_{d_{p-u_l}}} s_{i_{c_1}} s_{i_{c_2}} \cdots s_{i_{c_{u_l}}} \\ &= s_{i_{d_1}} s_{i_{d_2}} \cdots s_{i_{d_{p-u_l}}} s_{i_l} s_{i_{c_1}} \cdots s_{i_{c_{r-1}}} s_{i_{c_{r+1}}} \cdots s_{i_{c_{u_l}}} \end{aligned}$$

is a reduced expression of xw_0, and hence the sequence

$$(d_1,\, d_2,\, \ldots,\, d_{p-u_l},\, l,\, c_1,\, \ldots,\, c_{r-1},\, c_{r+1},\, \ldots,\, c_{u_l})$$

is an element of $S(xw_0,\, \mathbf{i})$. However, since $l < l + 1 \leq c_1$, this sequence is strictly less than $\min S(xw_0,\, \mathbf{i})$ with respect to the lexicographic ordering $\succeq$, which is a contradiction. Thus we have proved Lemma 4.1.3. □

4.3 Proof of Proposition 4.1.4. Keep the notation and assumptions of §4.1. Assume that $\mathbf{i} = (i_1, i_2, \ldots, i_m) \in R(w_0)$ and $\mathbf{j} = (j_1, j_2, \ldots, j_m) \in R(w_0)$ are related by a 2-move or a 3-move. We will study the relation between the two elements $\min S(xw_0, \mathbf{i})$ and $\min S(xw_0, \mathbf{j})$.

Assume first that $\mathbf{i}$ and $\mathbf{j}$ are related by a 2-move as in (2.1.3), i.e.,

$$\begin{aligned} \mathbf{i} &= (i_1, \ldots, i_k, i, j, i_{k+3}, \ldots, i_m), \\ \mathbf{j} &= (i_1, \ldots, i_k, j, i, i_{k+3}, \ldots, i_m) \end{aligned}$$

for some indices $i, j \in I$ with $a_{ij} = a_{ji} = 0$ and an integer $0 \le k \le m-2$. We define a map $\sigma_{\mathbf{i},\mathbf{j}} : S(xw_0, \mathbf{i}) \to S(xw_0, \mathbf{j})$ (resp., $\sigma_{\mathbf{j},\mathbf{i}} : S(xw_0, \mathbf{j}) \to S(xw_0, \mathbf{i})$) as follows. Let $\mathbf{a} = (a_1, a_2, \ldots, a_p) \in S(xw_0, \mathbf{i})$ (resp., $\mathbf{a} = (a_1, a_2, \ldots, a_p) \in S(xw_0, \mathbf{j})$). We have

$$\begin{aligned} &[1, k] \cap \mathbf{a} = (a_1, \ldots, a_{u_0}), \\ &[k+1, k+2] \cap \mathbf{a} = (a_{u_0+1}, \ldots, a_{u_1}), \\ &[k+3, m] \cap \mathbf{a} = (a_{u_1+1}, \ldots, a_p) \end{aligned}$$

for $0 \le u_0 \le u_1 \le p$. Then we set $\mathbf{a}' := [k+1, k+2] \cap \mathbf{a} = (a_{u_0+1}, \ldots, a_{u_1})$, and define a strictly increasing sequence $\mathbf{b} = (b_1, b_2, \ldots, b_p)$ by:

$$b_q = a_q \quad \text{for } 1 \le q \le u_0 \text{ and } u_1 + 1 \le q \le p,$$

$$(b_{u_0+1}, \ldots, b_{u_1}) = \begin{cases} \emptyset & \text{if } \mathbf{a}' = \emptyset, \\ (k+1) & \text{if } \mathbf{a}' = (k+2), \\ (k+2) & \text{if } \mathbf{a}' = (k+1), \\ (k+1, k+2) & \text{if } \mathbf{a}' = (k+1, k+2). \end{cases}$$

It is obvious that $\mathbf{b} \in S(xw_0, \mathbf{j})$ (resp., $\mathbf{b} \in S(xw_0, \mathbf{i})$). We now set $\sigma_{\mathbf{i},\mathbf{j}}(\mathbf{a}) := \mathbf{b}$ (resp., $\sigma_{\mathbf{j},\mathbf{i}}(\mathbf{a}) := \mathbf{b}$).

Remark 4.3.1. Assume that $\mathbf{i}$ and $\mathbf{j}$ are related by a 2-move as above. Obviously, $\sigma_{\mathbf{j},\mathbf{i}} \circ \sigma_{\mathbf{i},\mathbf{j}} = \mathrm{id}_{S(xw_0,\mathbf{i})}$ and $\sigma_{\mathbf{i},\mathbf{j}} \circ \sigma_{\mathbf{j},\mathbf{i}} = \mathrm{id}_{S(xw_0,\mathbf{j})}$.

Assume next that $\mathbf{i}$ and $\mathbf{j}$ are related by a 3-move as in (2.1.4), i.e.,

$$\begin{aligned} \mathbf{i} &= (i_1, \ldots, i_k, i, j, i, i_{k+4}, \ldots, i_m), \\ \mathbf{j} &= (i_1, \ldots, i_k, j, i, j, i_{k+4}, \ldots, i_m) \end{aligned}$$

for some indices $i, j \in I$ with $a_{ij} = a_{ji} = -1$ and an integer $0 \le k \le m-3$. We define a map $\sigma_{\mathbf{i},\mathbf{j}} : S(xw_0, \mathbf{i}) \to S(xw_0, \mathbf{j})$ (resp., $\sigma_{\mathbf{j},\mathbf{i}} : S(xw_0, \mathbf{j}) \to S(xw_0, \mathbf{i})$) as follows. Let $\mathbf{a} = (a_1, a_2, \ldots, a_p) \in S(xw_0, \mathbf{i})$ (resp., $\mathbf{a} = (a_1, a_2, \ldots, a_p) \in S(xw_0, \mathbf{j})$). We have

$$\begin{aligned} &[1, k] \cap \mathbf{a} = (a_1, \ldots, a_{u_0}), \\ &[k+1, k+3] \cap \mathbf{a} = (a_{u_0+1}, \ldots, a_{u_1}), \\ &[k+4, m] \cap \mathbf{a} = (a_{u_1+1}, \ldots, a_p) \end{aligned}$$

for $0 \le u_0 \le u_1 \le p$. Then we set $\mathbf{a}' := [k+1,\, k+3] \cap \mathbf{a} = (a_{u_0+1},\, \ldots,\, a_{u_1})$, and define a strictly increasing sequence $\mathbf{b} = (b_1,\, b_2,\, \ldots,\, b_p)$ by:

$$b_q = a_q \quad \text{for } 1 \le q \le u_0 \text{ and } u_1 + 1 \le q \le p,$$

$$(b_{u_0+1},\, \ldots,\, b_{u_1}) = \begin{cases} \emptyset & \text{if } \mathbf{a}' = \emptyset, \\ (k+2) & \text{if } \mathbf{a}' = (k+1) \text{ or } (k+3), \\ (k+1) & \text{if } \mathbf{a}' = (k+2), \\ (k+2,\, k+3) & \text{if } \mathbf{a}' = (k+1,\, k+2), \\ (k+1,\, k+2) & \text{if } \mathbf{a}' = (k+2,\, k+3), \\ (k+1,\, k+2,\, k+3) & \text{if } \mathbf{a}' = (k+1,\, k+2,\, k+3). \end{cases}$$

It is easily seen that $\mathbf{b} \in S(xw_0,\, \mathbf{j})$ (resp., $\mathbf{b} \in S(xw_0,\, \mathbf{i})$). We now set $\sigma_{\mathbf{i},\mathbf{j}}(\mathbf{a}) := \mathbf{b}$ (resp., $\sigma_{\mathbf{j},\mathbf{i}}(\mathbf{a}) := \mathbf{b}$).

Remark 4.3.2. Assume that $\mathbf{i}$ and $\mathbf{j}$ are related by a 3-move as above, and take $\mathbf{a} = (a_1,\, a_2,\, \ldots,\, a_p)$ to be the minimum element $\min S(xw_0,\, \mathbf{i})$. It follows from the minimality of $\mathbf{a} = (a_1,\, a_2,\, \ldots,\, a_p) \in S(xw_0,\, \mathbf{i})$ with respect to the lexicographic ordering that if $u_1 = u_0 + 1$, then $\mathbf{a}' = (k+1)$ or $(k+2)$. Using this, we see easily from the definitions of the maps $\sigma_{\mathbf{i},\mathbf{j}}$ and $\sigma_{\mathbf{j},\mathbf{i}}$ that

$$\sigma_{\mathbf{j},\mathbf{i}}\bigl(\sigma_{\mathbf{i},\mathbf{j}}(\min S(xw_0,\, \mathbf{i}))\bigr) = \min S(xw_0,\, \mathbf{i}).$$

Similarly, we obtain

$$\sigma_{\mathbf{i},\mathbf{j}}\bigl(\sigma_{\mathbf{j},\mathbf{i}}(\min S(xw_0,\, \mathbf{j}))\bigr) = \min S(xw_0,\, \mathbf{j}).$$

Proposition 4.3.3. *Assume that* $\mathbf{i} = (i_1,\, i_2,\, \ldots,\, i_m)$, $\mathbf{j} = (j_1,\, j_2,\, \ldots,\, j_m) \in R(w_0)$ *are related by a* 2*-move or a* 3*-move as above. Then, we have*

$$\sigma_{\mathbf{i},\mathbf{j}}(\min S(xw_0,\, \mathbf{i})) = \min S(xw_0,\, \mathbf{j}) \quad \textit{and} \quad \sigma_{\mathbf{j},\mathbf{i}}(\min S(xw_0,\, \mathbf{j})) = \min S(xw_0,\, \mathbf{i}).$$

Proof. We prove that $\sigma_{\mathbf{i},\mathbf{j}}(\min S(xw_0,\, \mathbf{i})) = \min S(xw_0,\, \mathbf{j})$ in the case that $\mathbf{i}$ and $\mathbf{j}$ are related by a 3-move as in (2.1.4); the proof of the equation $\sigma_{\mathbf{j},\mathbf{i}}(\min S(xw_0,\, \mathbf{j})) = \min S(xw_0,\, \mathbf{j})$ is similar, and the proof for the case of 2-move is simpler. For simplicity of notation, we set

$$\begin{aligned} &\mathbf{a} = (a_1,\, a_2,\, \ldots,\, a_p) := \min S(xw_0,\, \mathbf{i}), \\ &\mathbf{b} = (b_1,\, b_2,\, \ldots,\, b_p) := \sigma_{\mathbf{i},\mathbf{j}}(\min S(xw_0,\, \mathbf{i})) = \sigma_{\mathbf{i},\mathbf{j}}(\mathbf{a}), \\ &\mathbf{c} = (c_1,\, c_2,\, \ldots,\, c_p) := \min S(xw_0,\, \mathbf{j}), \\ &\mathbf{d} = (d_1,\, d_2,\, \ldots,\, d_p) := \sigma_{\mathbf{j},\mathbf{i}}(\min S(xw_0,\, \mathbf{j})) = \sigma_{\mathbf{j},\mathbf{i}}(\mathbf{c}). \end{aligned}$$

We have

$$\begin{cases} [1,\, k] \cap \mathbf{a} = (a_1,\, \ldots,\, a_{u_0}), \\ [k+1,\, k+3] \cap \mathbf{a} = (a_{u_0+1},\, \ldots,\, a_{u_1}), \\ [k+4,\, m] \cap \mathbf{a} = (a_{u_1+1},\, \ldots,\, a_p) \end{cases} \tag{4.3.1}$$

for $0 \le u_0 \le u_1 \le p$. From the definition of the map $\sigma_{\mathbf{i},\mathbf{j}}$, it is obvious that

$$\begin{aligned} & b_q = a_q \quad \text{for all } 1 \le q \le u_0 \text{ and } u_1 + 1 \le q \le p, \\ & [k+1,\, k+3] \cap \mathbf{b} = (b_{u_0+1},\, \dots,\, b_{u_1}). \end{aligned} \tag{4.3.2}$$

Also, we have

$$\begin{cases} [1,\, k] \cap \mathbf{c} = (c_1,\, \dots,\, c_{t_0}), \\ [k+1,\, k+3] \cap \mathbf{c} = (c_{t_0+1},\, \dots,\, c_{t_1}), \\ [k+4,\, m] \cap \mathbf{c} = (c_{t_1+1},\, \dots,\, c_p) \end{cases} \tag{4.3.3}$$

for $0 \le t_0 \le t_1 \le p$. From the definition of the map $\sigma_{\mathbf{j},\mathbf{i}}$, it is obvious that

$$\begin{aligned} & d_q = c_q \quad \text{for all } 1 \le q \le t_0 \text{ and } t_1 + 1 \le q \le p, \\ & [k+1,\, k+3] \cap \mathbf{d} = (d_{t_0+1},\, \dots,\, d_{t_1}). \end{aligned} \tag{4.3.4}$$

We will prove that $\mathbf{b} = \mathbf{c}$.

Since $\mathbf{b} = \sigma_{\mathbf{i},\mathbf{j}}(\mathbf{a}) \in S(xw_0,\, \mathbf{j})$, it follows that $\mathbf{b} \succeq \mathbf{c} = \min S(xw_0,\, \mathbf{j})$. Suppose, contrary to our claim, that $\mathbf{b} \succneqq \mathbf{c}$. Then, there exists an integer $1 \le q_0 \le p$ such that $b_q = c_q$ for all $1 \le q < q_0$ and $b_{q_0} > c_{q_0}$. If $q_0 \le \min\{u_0,\, t_0\}$, then it would follow from (4.3.2) and (4.3.4) that $a_q = d_q$ for all $1 \le q < q_0$ and $a_{q_0} > d_{q_0}$. This means that $\mathbf{a} \succneqq \mathbf{d}$, contradicting the minimality of $\mathbf{a} = \min S(xw_0,\, \mathbf{i})$. Therefore, we obtain $q_0 > \min\{u_0,\, t_0\}$, and hence $b_q = c_q$ for all $1 \le q \le \min\{u_0,\, t_0\}$. We now show that $u_0 = t_0$. Indeed, if $u_0 > t_0$, then we would have $b_{t_0+1} \le b_{u_0} \le k$ by (4.3.1) and (4.3.2). Since $c_{t_0+1} \ge k+1$ by (4.3.3), it follows that $b_{t_0+1} < c_{t_0+1}$. However, since $b_q = c_q$ for all $1 \le q \le \min\{u_0,\, t_0\} = t_0 < q_0$ as shown above, this implies that $\mathbf{b} \precneqq \mathbf{c} = \min S(xw_0,\, \mathbf{j})$, a contradiction. Similarly, if $u_0 < t_0$, then we would have $d_{u_0+1} \le d_{t_0} \le k$ by (4.3.3) and (4.3.4). Since $a_{u_0+1} \ge k+1$ by (4.3.1), it follows that $d_{u_0+1} < a_{u_0+1}$. Also, since $b_q = c_q$ for all $1 \le q \le \min\{u_0,\, t_0\} = u_0$, it follows from (4.3.2) and (4.3.4) that $d_q = a_q$ for all $1 \le q \le u_0$. Combining these, we obtain $\mathbf{d} \precneqq \mathbf{a} = \min S(xw_0,\, \mathbf{i})$, a contradiction. Hence we conclude that $u_0 = t_0$ and $b_q = c_q$ for all $1 \le q \le u_0 = t_0$; in addition, we see from (4.3.2) and (4.3.4) that $a_q = d_q$ for all $1 \le q \le u_0 = t_0$. Thus, we have shown that

$$\begin{aligned} & [1,\, k] \cap \mathbf{b} = (c_1,\, \dots,\, c_{t_0}) = [1,\, k] \cap \mathbf{c}, \\ & [k+1,\, k+3] \cap \mathbf{b} = (b_{t_0+1},\, \dots,\, b_{u_1}). \end{aligned}$$

Next, let us show that $[k+1,\, k+3] \cap \mathbf{b} = [k+1,\, k+3] \cap \mathbf{c}$. Here we give a proof in the case that $[k+1,\, k+3] \cap \mathbf{c} = (k+1,\, k+2)$; the proofs for the other cases are similar. Since $[1,\, k] \cap \mathbf{b} = [1,\, k] \cap \mathbf{c}$ as shown above and $\mathbf{b} \succneqq \mathbf{c}$ by assumption, the sequence $[k+1,\, k+3] \cap \mathbf{b}$ is not equal to $(k+1,\, k+2,\, k+3)$. Also, since $\mathbf{b} = \sigma_{\mathbf{i},\mathbf{j}}(\mathbf{a})$, it follows from the definition of the map $\sigma_{\mathbf{i},\mathbf{j}}$ that the sequence $[k+1,\, k+3] \cap \mathbf{b}$ cannot be equal to $(k+3)$ nor to $(k+1,\, k+3)$. Furthermore, if $[k+1,\, k+3] \cap \mathbf{b} = \emptyset$, then we would have

$t_0 = u_1$ and $a_{t_0+1} = a_{u_1+1} \geq k+4$. Since $a_q = d_q$ for all $1 \leq q \leq t_0$ as seen above, and since $d_{t_0+1} = k+2$ $(< a_{t_0+1})$ by the definition of the map $\sigma_{\mathbf{j},\mathbf{i}}$, along with the assumption that $[k+1, k+3] \cap \mathbf{c} = (k+1, k+2)$, it follows that $\mathbf{a} \gneqq \mathbf{d} = \sigma_{\mathbf{j},\mathbf{i}}(\mathbf{c})$, which contradicts the minimality of $\mathbf{a} = \min S(xw_0, \mathbf{i})$. Similarly, if $[k+1, k+3] \cap \mathbf{b} = (k+1)$, then we would have $t_0 + 1 = u_1$ and $a_{t_0+2} = a_{u_1+1} \geq k+4$; also, since $\sigma_{\mathbf{j},\mathbf{i}}(\mathbf{b}) = \mathbf{a}$ by Remark 4.3.2, it follows from the definition of the map $\sigma_{\mathbf{j},\mathbf{i}}$ that $a_{t_0+1} = a_{u_1} = k+2$. Since $a_q = d_q$ for all $1 \leq q \leq t_0$ as seen above, and since $d_{t_0+1} = k+2$ $(= a_{t_0+1})$ and $d_{t_0+2} = k+3$ $(< a_{t_0+2})$ by the definition of the map $\sigma_{\mathbf{j},\mathbf{i}}$ along with the assumption that $[k+1, k+3] \cap \mathbf{c} = (k+1, k+2)$, it follows that $\mathbf{a} \gneqq \mathbf{d} = \sigma_{\mathbf{j},\mathbf{i}}(\mathbf{c})$, contradicting the minimality of $\mathbf{a}$. Now, suppose, contrary to our claim, that $[k+1, k+3] \cap \mathbf{b} = (k+2)$ or $(k+2, k+3)$; note that $b_{t_0+1} = k+2$. We set $w' := s_{j_{b_1}} s_{j_{b_2}} \cdots s_{j_{b_{t_0}}} = s_{j_{c_1}} s_{j_{c_2}} \cdots s_{j_{c_{t_0}}}$, and $w := (w')^{-1} x w_0$. Then, the w has reduced expressions of the form

$$w = s_{j_{b_{t_0+1}}} s_{j_{b_{t_0+2}}} \cdots s_{j_{b_p}} = s_{j_{c_{t_0+1}}} s_{j_{c_{t_0+2}}} \cdots s_{j_{c_p}}.$$

If we set $z := s_{j_{c_{t_1+1}}} s_{j_{c_{t_1+2}}} \cdots s_{j_{c_p}}$, then

$$w = (s_{j_{c_{t_0+1}}} \ldots s_{j_{c_{t_1}}}) z = s_{j_{k+1}} s_{j_{k+2}} z = s_j s_i z$$

since $(c_{t_0+1}, \ldots, c_{t_1}) = (k+1, k+2)$ by assumption and $j_{k+1} = j$, $j_{k+2} = i$. Also, since $b_{t_0+1} = k+2$ by assumption and $j_{k+2} = i$, we deduce by [MoP, Chap. 5, §3, Proposition 2 (i)] that $w^{-1} \cdot \alpha_{j_{k+2}} = w^{-1} \cdot \alpha_i$ is a negative root, and so $z^{-1} \cdot \alpha_j = z^{-1} s_i s_j \cdot \alpha_i = (s_j s_i z)^{-1} \cdot \alpha_i = w^{-1} \cdot \alpha_i$ is a negative root. Hence it follows from the exchange condition that z has a reduced expression of the form:

$$z = s_j s_{j_{c_{t_1+1}}} \cdots s_{j_{c_{r-1}}} s_{j_{c_{r+1}}} \cdots s_{j_{c_p}}$$

for some $t_1 + 1 \leq r \leq p$. Therefore, we obtain

$$\begin{aligned} xw_0 &= w'w = w' s_j s_i z \\ &= s_{j_{c_1}} s_{j_{c_2}} \cdots s_{j_{c_{t_0}}} s_j s_i s_j s_{j_{c_{t_1+1}}} \cdots s_{j_{c_{r-1}}} s_{j_{c_{r+1}}} \cdots s_{j_{c_p}} \\ &= s_{j_{c_1}} s_{j_{c_2}} \cdots s_{j_{c_{t_0}}} s_{j_{k+1}} s_{j_{k+2}} s_{j_{k+3}} s_{j_{c_{t_1+1}}} \cdots s_{j_{c_{r-1}}} s_{j_{c_{r+1}}} \cdots s_{j_{c_p}}, \end{aligned}$$

which is a reduced expression of xw_0, and hence the sequence

$$(c_1, c_2, \ldots, c_{t_0}, k+1, k+2, k+3, c_{t_1+1}, \ldots, c_{r-1}, c_{r+1}, \ldots, c_p)$$

is an element of $S(xw_0, \mathbf{j})$. However, this sequence is strictly less than $\mathbf{c} = \min S(xw_0, \mathbf{j})$ with respect to the lexicographic ordering, which is a contradiction. Hence we conclude that $[k+1, k+3] \cap \mathbf{b} = (k+1, k+2) = [k+1, k+3] \cap \mathbf{c}$. Thus, we have shown that $u_1 = t_1$ and $b_q = c_q$ for $t_0 + 1 \leq q \leq u_1 = t_1$, and so

$$\begin{cases} [1, k] \cap \mathbf{b} = (c_1, \ldots, c_{t_0}) = [1, k] \cap \mathbf{c}, \\ [k+1, k+3] \cap \mathbf{b} = (c_{t_0+1}, \ldots, c_{t_1}) = [k+1, k+3] \cap \mathbf{c}, \\ [k+4, m] \cap \mathbf{b} = (b_{t_1+1}, \ldots, b_p). \end{cases} \tag{4.3.5}$$

The equations (4.3.5) imply that $q_0 > t_1 = u_1$. In this case, since $\mathbf{c} = \min S(xw_0, \mathbf{j}) \precneqq \mathbf{b}$, it follows from the definition of the map $\sigma_{\mathbf{j},\mathbf{i}}$ that $\sigma_{\mathbf{j},\mathbf{i}}(\mathbf{c}) \precneqq \sigma_{\mathbf{j},\mathbf{i}}(\mathbf{b}) = \mathbf{a}$, contrary to the minimality of $\mathbf{a}$. Thus we have proved that $\mathbf{b} = \mathbf{c}$, thereby completing the proof of Proposition 4.3.3. $\square$

Proof of Proposition 4.1.4. By Remark 2.1.2, we may assume that $\mathbf{i}, \mathbf{j} \in R(w_0)$ are related by a 2-move or a 3-move. Furthermore, in view of [Kam1, Theorem 7.1], it suffices to show that $\mu^{x,\mathbf{i}}_{w^{\mathbf{j}}_l} = \xi^{\mathbf{j}}_l$ for every $0 \le l \le m$. We give a proof in the case that $\mathbf{i}, \mathbf{j} \in R(w_0)$ are related by a 3-move as in (2.1.4), i.e.,

$$\begin{aligned} \mathbf{i} &= (i_1, \dots, i_k, i, j, i, i_{k+4}, \dots, i_m), \\ \mathbf{j} &= (i_1, \dots, i_k, j, i, j, i_{k+4}, \dots, i_m) \end{aligned}$$

for some indices $i, j \in I$ with $a_{ij} = a_{ji} = -1$ and an integer $0 \le k \le m-3$; the proof for the case of 2-move is similar (or, even simpler).

Assume first that $l \ne k+1, k+2$. Then, it is obvious that $w^{\mathbf{i}}_l = w^{\mathbf{j}}_l$. Also, because $\min S(xw_0, \mathbf{j}) = \sigma_{\mathbf{i},\mathbf{j}}(\min S(xw_0, \mathbf{i}))$ by Proposition 4.3.3, we deduce from the definition (see §4.2) that $v^{\mathbf{i}}_l = v^{\mathbf{j}}_l$. Therefore, it follows immediately from Lemma 4.2.1 that

$$\xi^{\mathbf{i}}_l = w^{\mathbf{i}}_l v^{\mathbf{i}}_l w_0^{-1} \cdot \lambda = w^{\mathbf{j}}_l v^{\mathbf{j}}_l w_0^{-1} \cdot \lambda = \xi^{\mathbf{j}}_l.$$

Hence we obtain $\mu^{x,\mathbf{i}}_{w^{\mathbf{j}}_l} = \mu^{x,\mathbf{i}}_{w^{\mathbf{i}}_l} = \xi^{\mathbf{i}}_l = \xi^{\mathbf{j}}_l$, as desired. Thus, it remains to show that

$$\mu^{x,\mathbf{i}}_{w^{\mathbf{j}}_{k+1}} = \xi^{\mathbf{j}}_{k+1} \quad \text{and} \quad \mu^{x,\mathbf{i}}_{w^{\mathbf{j}}_{k+2}} = \xi^{\mathbf{j}}_{k+2}. \tag{4.3.6}$$

Recall that the integers $n^{\mathbf{i}}_l = n^{\mathbf{i}}_l(\mu^{x,\mathbf{i}}_\bullet) \in \mathbb{Z}_{\ge 0}$ and $n^{\mathbf{j}}_l = n^{\mathbf{j}}_l(\mu^{x,\mathbf{i}}_\bullet) \in \mathbb{Z}_{\ge 0}$ for $l = k+1, k+2, k+3$ are defined via the length formula:

$$\mu^{x,\mathbf{i}}_{w^{\mathbf{i}}_l} - \mu^{x,\mathbf{i}}_{w^{\mathbf{i}}_{l-1}} = n^{\mathbf{i}}_l(w^{\mathbf{i}}_{l-1} \cdot h_{i_l}) \quad \text{and} \quad \mu^{x,\mathbf{i}}_{w^{\mathbf{j}}_l} - \mu^{x,\mathbf{i}}_{w^{\mathbf{j}}_{l-1}} = n^{\mathbf{j}}_l(w^{\mathbf{j}}_{l-1} \cdot h_{j_l}),$$

respectively. Also, we know from (the proof of) Lemma 4.1.3 that for $l = k+1, k+2, k+3$, there exists an integer $N^{\mathbf{j}}_l \in \mathbb{Z}_{\ge 0}$ such that

$$\xi^{\mathbf{j}}_l - \xi^{\mathbf{j}}_{l-1} = N^{\mathbf{j}}_l(w^{\mathbf{j}}_{l-1} \cdot h_{j_l});$$

recall that the integer $N^{\mathbf{j}}_l \in \mathbb{Z}_{\ge 0}$ is given by:

$$N^{\mathbf{j}}_l = \begin{cases} 0 & \text{if } l \text{ appears in } \min S(xw_0, \mathbf{j}), \\ \langle \beta^{\mathbf{j}}_l, \xi^{\mathbf{j}}_l \rangle & \text{otherwise.} \end{cases} \tag{4.3.7}$$

Since $\mu^{x,\mathbf{i}}_{w^{\mathbf{j}}_{k+3}} = \xi^{\mathbf{j}}_{k+3}$ as shown above, equations (4.3.6) will follow if we can show that $n^{\mathbf{j}}_{k+3} = N^{\mathbf{j}}_{k+3}$ and $n^{\mathbf{j}}_{k+2} = N^{\mathbf{j}}_{k+2}$. We will show that $n^{\mathbf{j}}_l = N^{\mathbf{j}}_l$ for $l = k+1, k+2, k+3$.

We know that the sequence $[k+1,\, k+3] \cap \min S(xw_0,\, \mathbf{i})$ is equal to one of the following: (i) $\emptyset$, (ii) $(k+1)$, (iii) $(k+2)$, (iv) $(k+1,\, k+2)$, (v) $(k+2,\, k+3)$, (vi) $(k+1,\, k+2,\, k+3)$. Also, since $\mu^{x,\mathbf{i}}_{w^{\mathbf{i}}_l} = \xi^{\mathbf{i}}_l$ for all $0 \le l \le m$ and hence $\mu^{x,\mathbf{i}}_{w^{\mathbf{i}}_l} - \mu^{x,\mathbf{i}}_{w^{\mathbf{i}}_{l-1}} = \xi^{\mathbf{i}}_l - \xi^{\mathbf{i}}_{l-1}$ for all $1 \le l \le m$, it follows from (4.2.2) that

$$n^{\mathbf{i}}_{k+1} = \begin{cases} 0 & \text{if } k+1 \text{ appears in } \min S(xw_0,\, \mathbf{i}), \\ \langle \beta^{\mathbf{i}}_{k+1},\, \xi^{\mathbf{i}}_{k+1} \rangle & \text{otherwise}, \end{cases} \tag{4.3.8}$$

$$n^{\mathbf{i}}_{k+2} = \begin{cases} 0 & \text{if } k+2 \text{ appears in } \min S(xw_0,\, \mathbf{i}), \\ \langle \beta^{\mathbf{i}}_{k+2},\, \xi^{\mathbf{i}}_{k+2} \rangle & \text{otherwise}, \end{cases} \tag{4.3.9}$$

$$n^{\mathbf{i}}_{k+3} = \begin{cases} 0 & \text{if } k+3 \text{ appears in } \min S(xw_0,\, \mathbf{i}), \\ \langle \beta^{\mathbf{i}}_{k+3},\, \xi^{\mathbf{i}}_{k+3} \rangle & \text{otherwise}. \end{cases} \tag{4.3.10}$$

For simplicity of notation, we set $\gamma_i := w^{\mathbf{i}}_k \cdot \alpha_i$ and $\gamma_j := w^{\mathbf{i}}_k \cdot \alpha_j$. Then we calculate:

$$\begin{aligned} \beta^{\mathbf{i}}_{k+1} &= w^{\mathbf{i}}_k \cdot \alpha_{i_{k+1}} = w^{\mathbf{i}}_k \cdot \alpha_i = \gamma_i, \\ \beta^{\mathbf{i}}_{k+2} &= w^{\mathbf{i}}_{k+1} \cdot \alpha_{i_{k+2}} = w^{\mathbf{i}}_k s_{i_{k+1}} \cdot \alpha_{i_{k+2}} = w^{\mathbf{i}}_k s_i \cdot \alpha_j = w^{\mathbf{i}}_k \cdot (\alpha_i + \alpha_j) = \gamma_i + \gamma_j, \\ \beta^{\mathbf{i}}_{k+3} &= w^{\mathbf{i}}_{k+2} \cdot \alpha_{i_{k+3}} = w^{\mathbf{i}}_k s_{i_{k+1}} s_{i_{k+2}} \cdot \alpha_{i_{k+3}} = w^{\mathbf{i}}_k s_i s_j \cdot \alpha_i = w^{\mathbf{i}}_k \cdot \alpha_j = \gamma_j. \end{aligned}$$

Also, we set $A := \langle \gamma_i,\, \xi^{\mathbf{i}}_{k+3} \rangle$ and $B := \langle \gamma_j,\, \xi^{\mathbf{i}}_{k+3} \rangle$. Then, we obtain

$$(n^{\mathbf{i}}_{k+1},\, n^{\mathbf{i}}_{k+2},\, n^{\mathbf{i}}_{k+3}) = \begin{cases} (B,\, A,\, B) & \text{in case (i)}, \\ (0,\, A,\, B) & \text{in case (ii)}, \\ (A+B,\, 0,\, B) & \text{in case (iii)}, \\ (0,\, 0,\, B) & \text{in case (iv)}, \\ (A,\, 0,\, 0) & \text{in case (v)}, \\ (0,\, 0,\, 0) & \text{in case (vi)}. \end{cases} \tag{4.3.11}$$

As an example, let us give a proof in case (iii). Since $k+3$ does not appear in $\min S(xw_0,\, \mathbf{i})$, we have $\xi^{\mathbf{i}}_{k+2} = s_{\beta^{\mathbf{i}}_{k+3}} \cdot \xi^{\mathbf{i}}_{k+3}$ by definition (4.1.1), and $n^{\mathbf{i}}_{k+3} = \langle \beta^{\mathbf{i}}_{k+3},\, \xi^{\mathbf{i}}_{k+3} \rangle = \langle \gamma_j,\, \xi^{\mathbf{i}}_{k+3} \rangle = B$ by (4.3.10). Since $k+2$ appears in $\min S(xw_0,\, \mathbf{i})$, we have $\xi^{\mathbf{i}}_{k+1} = \xi^{\mathbf{i}}_{k+2} = s_{\beta^{\mathbf{i}}_{k+3}} \cdot \xi^{\mathbf{i}}_{k+3}$ by definition (4.1.1), and $n^{\mathbf{i}}_{k+2} = 0$ by (4.3.9). Since $k+1$ does not appear in $\min S(xw_0,\, \mathbf{i})$, we conclude by (4.3.8) that

$$\begin{aligned} n^{\mathbf{i}}_{k+1} &= \langle \beta^{\mathbf{i}}_{k+1},\, \xi^{\mathbf{i}}_{k+1} \rangle = \langle \beta^{\mathbf{i}}_{k+1},\, s_{\beta^{\mathbf{i}}_{k+3}} \cdot \xi^{\mathbf{i}}_{k+3} \rangle = \langle \gamma_i,\, s_{\gamma_j} \cdot \xi^{\mathbf{i}}_{k+3} \rangle \\ &= \langle \gamma_i + \gamma_j,\, \xi^{\mathbf{i}}_{k+3} \rangle = A + B. \end{aligned}$$

The proofs for the other cases are similar. Hence it follows from (2.1.6) that

$$(n^{\mathbf{j}}_{k+1},\, n^{\mathbf{j}}_{k+2},\, n^{\mathbf{j}}_{k+3}) = \begin{cases} (A,\, B,\, A) & \text{in case (i)}, \\ (A+B,\, 0,\, A) & \text{in case (ii)}, \\ (0,\, B,\, A) & \text{in case (iii)}, \\ (B,\, 0,\, 0) & \text{in case (iv)}, \\ (0,\, 0,\, A) & \text{in case (v)}, \\ (0,\, 0,\, 0) & \text{in case (vi)}. \end{cases} \tag{4.3.12}$$

Next, we compute $(N^{\mathbf{j}}_{k+1},\, N^{\mathbf{j}}_{k+2},\, N^{\mathbf{j}}_{k+3})$. From Proposition 4.3.3, we deduce by the definition of the map $\sigma_{\mathbf{i},\mathbf{j}}$ that

$$[k+1,\, k+3] \cap \min S(xw_0,\, \mathbf{j}) = \begin{cases} \emptyset & \text{in case (i)}, \\ (k+2) & \text{in case (ii)}, \\ (k+1) & \text{in case (iii)}, \\ (k+2,\, k+3) & \text{in case (iv)}, \\ (k+1,\, k+2) & \text{in case (v)}, \\ (k+1,\, k+2,\, k+3) & \text{in case (vi)}. \end{cases}$$

Also, since $w^{\mathbf{i}}_k = w^{\mathbf{j}}_k$ and $\xi^{\mathbf{i}}_{k+3} = \xi^{\mathbf{j}}_{k+3}$ as seen above, we have $\gamma_i = w^{\mathbf{j}}_k \cdot \alpha_i$, $\gamma_j = w^{\mathbf{j}}_k \cdot \alpha_j$, and also $A = \langle \gamma_i,\, \xi^{\mathbf{j}}_{k+3} \rangle$, $B = \langle \gamma_j,\, \xi^{\mathbf{j}}_{k+3} \rangle$. Using these, by an argument similar to the one for (4.3.11), we obtain

$$(N^{\mathbf{j}}_{k+1},\, N^{\mathbf{j}}_{k+2},\, N^{\mathbf{j}}_{k+3}) = \begin{cases} (A,\, B,\, A) & \text{in case (i)}, \\ (A+B,\, 0,\, A) & \text{in case (ii)}, \\ (0,\, B,\, A) & \text{in case (iii)}, \\ (B,\, 0,\, 0) & \text{in case (iv)}, \\ (0,\, 0,\, A) & \text{in case (v)}, \\ (0,\, 0,\, 0) & \text{in case (vi)}. \end{cases} \tag{4.3.13}$$

As an example, let us give a proof in case (iii); recall that $[k+1,\, k+3] \cap \min S(xw_0,\, \mathbf{j}) = (k+1)$. Since $k+3$ does not appear in $\min S(xw_0,\, \mathbf{j})$, we have $\xi^{\mathbf{j}}_{k+2} = s_{\beta^{\mathbf{j}}_{k+3}} \cdot \xi^{\mathbf{j}}_{k+3}$ by definition (4.1.1), and $N^{\mathbf{j}}_{k+3} = \langle \beta^{\mathbf{j}}_{k+3},\, \xi^{\mathbf{j}}_{k+3} \rangle$ by (4.3.7). Also, we calculate:

$$\beta^{\mathbf{j}}_{k+3} = w^{\mathbf{j}}_{k+2} \cdot \alpha_{j_{k+3}} = w^{\mathbf{j}}_k s_{j_{k+1}} s_{j_{k+2}} \cdot \alpha_{j_{k+3}} = w^{\mathbf{j}}_k s_j s_i \cdot \alpha_j = w^{\mathbf{j}}_k \cdot \alpha_i = \gamma_i.$$

Hence we get $N^{\mathbf{j}}_{k+3} = \langle \gamma_i,\, \xi^{\mathbf{j}}_{k+3} \rangle = A$. Similarly, since $k+2$ does not appear in $\min S(xw_0,\, \mathbf{j})$, we have $N^{\mathbf{j}}_{k+2} = \langle \beta^{\mathbf{j}}_{k+2},\, \xi^{\mathbf{j}}_{k+2} \rangle$ and $\beta^{\mathbf{j}}_{k+2} = \gamma_i + \gamma_j$. Therefore, we conclude that

$$\begin{aligned} N^{\mathbf{j}}_{k+2} &= \langle \beta^{\mathbf{j}}_{k+2},\, \xi^{\mathbf{j}}_{k+2} \rangle = \langle \beta^{\mathbf{j}}_{k+2},\, s_{\beta^{\mathbf{j}}_{k+3}} \cdot \xi^{\mathbf{j}}_{k+3} \rangle \\ &= \langle \gamma_i + \gamma_j,\, s_{\gamma_i} \cdot \xi^{\mathbf{j}}_{k+3} \rangle = \langle \gamma_j,\, \xi^{\mathbf{j}}_{k+3} \rangle = B. \end{aligned}$$

Since $k+1$ appears in $\min S(xw_0,\, \mathbf{j})$, we have $\xi^{\mathbf{j}}_k = \xi^{\mathbf{j}}_{k+1}$ by definition (4.1.1), and hence $N^{\mathbf{j}}_{k+1} = 0$. The proofs for the other cases are similar.

Combining (4.3.12) and (4.3.13), we obtain $(n^{\mathbf{j}}_{k+1},\, n^{\mathbf{j}}_{k+2},\, n^{\mathbf{j}}_{k+3}) = (N^{\mathbf{j}}_{k+1},\, N^{\mathbf{j}}_{k+2},\, N^{\mathbf{j}}_{k+3})$, as desired. This completes the proof of Proposition 4.1.4. $\square$

4.4 Proof of Theorem 4.1.5. Keep the notation and assumptions of §4.1. Let $\mathbf{i} \in R(w_0)$, and write the element $\min S(xw_0,\, \mathbf{i})$ as:

$$\min S(xw_0,\, \mathbf{i}) = (a_1,\, a_2,\, \dots,\, a_p).$$

Then, we have

$$\begin{aligned} \mu^x_e = \mu^x_{w^{\mathbf{i}}_0} &= \xi^{\mathbf{i}}_0 = w^{\mathbf{i}}_0 v^{\mathbf{i}}_0 w_0^{-1} \cdot \lambda \quad \text{by (4.2.1)} \\ &= w^{\mathbf{i}}_0 \,(s_{i_{a_1}} s_{i_{a_2}} \cdots s_{i_{a_p}})\, w_0^{-1} \cdot \lambda = e(xw_0) w_0^{-1} \cdot \lambda = x \cdot \lambda. \end{aligned}$$

Therefore, it follows that $\mathrm{wt}(P(\mu^x_\bullet)) = \mu^x_e = x \cdot \lambda$. Because $P_{x\cdot\lambda}$ is the unique element of $\mathcal{MV}(\lambda)$ whose weight is $x \cdot \lambda$, we conclude that $P(\mu^x_\bullet) = P_{x\cdot\lambda}$. Thus, we have proved part (1) of Theorem 4.1.5.

Let us prove part (2). We first show that $P_{x\cdot\lambda} = P(\mu^x_\bullet) \subset \mathrm{Conv}(W_{\le x} \cdot \lambda)$. By (4.1.2), it suffices to show that $\xi^{\mathbf{i}}_l \in W_{\le x} \cdot \lambda$ for all $\mathbf{i} \in R(w_0)$ and $0 \le l \le m$. We take and fix an arbitrary $\mathbf{i} \in R(w_0)$, and show the assertion by induction on l. If $l = 0$, then the assertion is obvious since $\xi^{\mathbf{i}}_0 = x \cdot \lambda$ as shown above. Assume now that $l > 0$. If l appears in $\min S(xw_0,\, \mathbf{i})$, then $\xi^{\mathbf{i}}_l = \xi^{\mathbf{i}}_{l-1}$ by definition (4.1.1), and hence $\xi^{\mathbf{i}}_l \in W_{\le x} \cdot \lambda$ by our induction hypothesis. So, we may assume that l does not appear in $\min S(xw_0,\, \mathbf{i})$; note that $\xi^{\mathbf{i}}_{l-1} = s_{\beta^{\mathbf{i}}_l} \cdot \xi^{\mathbf{i}}_l$ by definition (4.1.1). In this case, we know from the proof of Lemma 4.1.3 that $\langle \beta^{\mathbf{i}}_l,\, \xi^{\mathbf{i}}_l \rangle \ge 0$. When $\langle \beta^{\mathbf{i}}_l,\, \xi^{\mathbf{i}}_l \rangle = 0$, we have $\xi^{\mathbf{i}}_l = \xi^{\mathbf{i}}_{l-1}$ by definition (see also (4.2.2)), and hence $\xi^{\mathbf{i}}_l \in W_{\le x} \cdot \lambda$ by our induction hypothesis. Now, it remains to consider the case $\langle \beta^{\mathbf{i}}_l,\, \xi^{\mathbf{i}}_l \rangle > 0$. By our induction hypothesis, there exists $z \in W_{\le x}$ such that $\xi^{\mathbf{i}}_{l-1} = z \cdot \lambda$. Then, we calculate:

$$\langle z^{-1} \cdot \beta^{\mathbf{i}}_l,\, \lambda \rangle = \langle \beta^{\mathbf{i}}_l,\, z \cdot \lambda \rangle = \langle \beta^{\mathbf{i}}_l,\, \xi^{\mathbf{i}}_{l-1} \rangle = \langle \beta^{\mathbf{i}}_l,\, s_{\beta^{\mathbf{i}}_l} \cdot \xi^{\mathbf{i}}_l \rangle = -\langle \beta^{\mathbf{i}}_l,\, \xi^{\mathbf{i}}_l \rangle < 0.$$

Since $\lambda \in X_*(T) \subset \mathfrak{h}_{\mathbb{R}}$ is a dominant coweight, this implies that $z^{-1} \cdot \beta^{\mathbf{i}}_l$ is a negative root. Therefore, we deduce from [MoP, Chap. 5, §3, Proposition 2 (i)] that $s_{\beta^{\mathbf{i}}_l} z < z \le x$, and hence that $\xi^{\mathbf{i}}_l = s_{\beta^{\mathbf{i}}_l} \cdot \xi^{\mathbf{i}}_{l-1} = s_{\beta^{\mathbf{i}}_l} z \cdot \lambda \in W_{\le x} \cdot \lambda$. Thus, we have shown that $P_{x\cdot\lambda} = P(\mu^x_\bullet) \subset \mathrm{Conv}(W_{\le x} \cdot \lambda)$ in all cases.

Next, we show the reverse inclusion: $P_{x\cdot\lambda} = P(\mu^x_\bullet) \supset \mathrm{Conv}(W_{\le x} \cdot \lambda)$. By (4.1.2), it suffices to show that for each $z \in W$ with $z \le x$, there exist some $\mathbf{i} \in R(w_0)$ and an integer $0 \le l \le m$ such that $\xi^{\mathbf{i}}_l = z \cdot \lambda$. Let $z \in W$ be such that $z \le x$; note that $zw_0 \ge xw_0$ (see, for example, [BjB, Proposition 2.3.4 (i)]). Denote by l the length $\ell(zw_0)$ of zw_0, and take $\mathbf{i} \in R(w_0)$ such that $w^{\mathbf{i}}_l = zw_0$. We define a subset $S(xw_0,\, \mathbf{i})_{\le l}$ of $S(xw_0,\, \mathbf{i})$ by:

$$S(xw_0,\, \mathbf{i})_{\le l} = \bigl\{ (a_1,\, a_2,\, \dots,\, a_p) \in S(xw_0,\, \mathbf{i}) \mid a_p \le l \bigr\} \subset S(xw_0,\, \mathbf{i});$$

since $w_l^{\mathbf{i}} = zw_0 \geq xw_0$, it follows from the "subword condition" (see, for example, [MoP, Chap. 5, §4, Proposition 2]) that $S(xw_0, \mathbf{i})_{\leq l}$ is not empty.

Claim. *The minimum element* $\min S(xw_0, \mathbf{i})_{\leq l}$ *of* $S(xw_0, \mathbf{i})_{\leq l}$ *with respect to the lexicographic ordering* $\succeq$ *is identical to the minimum element* $\min S(xw_0, \mathbf{i})$ *of* $S(xw_0, \mathbf{i})$*. In particular, we have* $[l+1, m] \cap \min S(xw_0, \mathbf{i}) = \emptyset$*.*

Proof of Claim. Let us write the elements $\min S(xw_0, \mathbf{i})_{\leq l}$ and $\min S(xw_0, \mathbf{i})$ as

$$\min S(xw_0, \mathbf{i})_{\leq l} = (a_1, a_2, \ldots, a_p) \quad \text{and} \quad \min S(xw_0, \mathbf{i}) = (b_1, b_2, \ldots, b_p),$$

respectively. Suppose, contrary to our claim, that $\min S(xw_0, \mathbf{i})_{\leq l} \gneqq \min S(xw_0, \mathbf{i})$. Then, there exists some integer $1 \leq q \leq p$ such that $a_1 = b_1, a_2 = b_2, \ldots, a_{q-1} = b_{q-1}$ and $a_q > b_q$; note that if $q \geq 2$, then $a_q > b_q > b_{q-1} = a_{q-1}$. We set $w := s_{i_{a_q}} \cdots s_{i_{a_p}}$. Since $s_{i_{a_1}} \cdots s_{i_{a_{q-1}}} = s_{i_{b_1}} \cdots s_{i_{b_{q-1}}}$ by the definition of q, and since

$$s_{i_{a_1}} \cdots s_{i_{a_{q-1}}} s_{i_{a_q}} \cdots s_{i_{a_p}} = s_{i_{b_1}} \cdots s_{i_{b_{q-1}}} s_{i_{b_q}} \cdots s_{i_{b_p}} = xw_0,$$

we get $w = s_{i_{b_q}} \cdots s_{i_{b_p}}$. It follows from [MoP, Chap. 5, §3, Proposition 2 (i)] that $w^{-1} \cdot \alpha_{i_{b_q}} = (s_{i_{b_q}} \cdots s_{i_{b_p}})^{-1} \cdot \alpha_{i_{b_q}}$ is a negative root. Therefore, we see by the exchange condition that w has a reduced expression of the form $s_{i_{b_q}} s_{i_{a_q}} \cdots s_{i_{a_{r-1}}} s_{i_{a_{r+1}}} \cdots s_{i_{a_p}}$ for some $q \leq r \leq p$, and hence that xw_0 has a reduced expression of the form

$$xw_0 = s_{i_{a_1}} \cdots s_{i_{a_{q-1}}} s_{i_{b_q}} s_{i_{a_q}} \cdots s_{i_{a_{r-1}}} s_{i_{a_{r+1}}} \cdots s_{i_{a_p}}.$$

It follows that the sequence

$$(a_1, \ldots, a_{q-1}, b_q, a_q, \ldots, a_{r-1}, a_{r+1}, \ldots, a_p)$$

is an element of $S(xw_0, \mathbf{i})_{\leq l}$, and is strictly less than $\mathbf{a} = \min S(xw_0, \mathbf{i})_{\leq l}$ with respect to the lexicographic ordering, which is a contradiction. Thus, we have shown that $\min S(xw_0, \mathbf{i})_{\leq l} = \min S(xw_0, \mathbf{i})$, as desired. This proves the claim. ▮

By (4.2.1), we have $\xi_l^{\mathbf{i}} = w_l^{\mathbf{i}} v_l^{\mathbf{i}} w_0^{-1} \cdot \lambda$. Since $[l+1, m] \cap \min S(xw_0, \mathbf{i}) = \emptyset$ by the claim above, it follows from the definition that $v_l^{\mathbf{i}} = e$. Hence we conclude that

$$\xi_l^{\mathbf{i}} = w_l^{\mathbf{i}} v_l^{\mathbf{i}} w_0^{-1} \cdot \lambda = w_l^{\mathbf{i}} w_0^{-1} \cdot \lambda = (zw_0) w_0^{-1} \cdot \lambda = z \cdot \lambda,$$

since $w_l^{\mathbf{i}} = zw_0$ for the $\mathbf{i} \in R(w_0)$. Thus, we have shown that $P_{x \cdot \lambda} = P(\mu_\bullet^x) \supset \mathrm{Conv}(W_{\leq x} \cdot \lambda)$, thereby completing the proof of part (2) of Theorem 4.1.5.

4.5 Relation between extremal MV polytopes and opposite Demazure crystals.

By combining Theorems 3.5.1 and 4.1.5, we obtain a polytopal condition for an MV polytope $P \in \mathcal{MV}(\lambda)$ to lie in the opposite Demazure crystal $\mathcal{MV}^{x}(\lambda)$ for $x \in W$.

Theorem 4.5.1. *Let* $x \in W$*. An element* $P \in \mathcal{MV}(\lambda)$ *is contained in* $\mathcal{MV}^{x}(\lambda) = \Psi_{\lambda}(\mathcal{B}^{x}(\lambda))$ *if and only if the MV polytope* P *contains* (*as a set*) *the extremal MV polytope* $P_{x\cdot\lambda} = \mathrm{Conv}(W_{\leq x} \cdot \lambda)$.

First, we prove the "only if" part of the theorem. Before giving the proof, we make the following remarks.

Remark 4.5.2. We know (see [BaG, §4.1] and [Kam2, Theorem 4.7]) that for $P \in \mathcal{MV}(\lambda)$ and $j \in I$ with $f_j P \neq \mathbf{0}$, there holds $f_j P \supset P$. Therefore, if an MV polytope $P \in \mathcal{MV}^{x}(\lambda)$ were obtained from the extremal MV polytope $P_{x\cdot\lambda}$ by successive application of the lowering Kashiwara operators f_j, $j \in I$, then it would follow immediately that the polytope P contains $P_{x\cdot\lambda}$. However, in general, not all MV polytopes in $\mathcal{MV}^{x}(\lambda)$ can be obtained from $P_{x\cdot\lambda}$ by successively applying f_j, $j \in I$, as Example 4.5.3 below shows.

Example 4.5.3. Let $\mathfrak{g}$ be the simple Lie algebra of type A_2. Set $\lambda := h_1 + h_2 \in X_*(T) \subset \mathfrak{h}_{\mathbb{R}}$, and $x := s_1 \in W$. It follows from Lemma 3.4.2 that an element $P \in \mathcal{MV}(\lambda)$ is contained in $\mathcal{MV}^{x}(\lambda)$ if and only if $f_1^{\max} f_2^{\max} P = P_{w_0\cdot\lambda}$. Hence we see from the crystal graph (4.5.1) below of $\mathcal{MV}(\lambda) \cong \mathcal{B}(\lambda)$ that $f_1^2 f_2 P_\lambda$ is contained in $\mathcal{MV}^{x}(\lambda)$. However, we deduce from the crystal graph (4.5.1) that $f_1^2 f_2 P_\lambda$ cannot be obtained from $P_{x\cdot\lambda}$ by successively applying f_1 and f_2.



(4.5.1)

Proof of the "only if" part of Theorem 4.5.1. Let $P = P(\mu_\bullet) \in \mathcal{MV}^{x}(\lambda)$ be an MV polytope with GGMS datum $\mu_\bullet = (\mu_w)_{w\in W}$. Since $P = P(\mu_\bullet) = \mathrm{Conv}\,\{\mu_w \mid w \in W\}$, and since $P_{x\cdot\lambda} = \mathrm{Conv}(W_{\leq x} \cdot \lambda)$ by Theorem 4.1.5, it suffices to show that for each $z \in W_{\leq x}$, there exists some $w \in W$ such that $\mu_w = z \cdot \lambda$. Now, let us take an arbitrary $z \in W_{\leq x}$, and set $q := \ell(zw_0)$. Then we know (see [Kas2, Proposition 3.2.4] and the comment following [Kas2, Proposition 4.3]) that $\mathcal{MV}^{z}(\lambda) \supset \mathcal{MV}^{x}(\lambda)$, which implies that $P \in \mathcal{MV}^{z}(\lambda)$. Therefore, there exists $\mathbf{i} = (i_1, i_2, \dots, i_m) \in R(w_0)$ with $w_q^{\mathbf{i}} = zw_0$ such that $\mu_{w_l^{\mathbf{i}}} = w_l^{\mathbf{i}} w_0 \cdot \lambda$ for all $q \le l \le m$. In particular, $\mu_{w_q^{\mathbf{i}}} = w_q^{\mathbf{i}} w_0 \cdot \lambda = (zw_0)w_0 \cdot \lambda = z \cdot \lambda$. Thus, we have proved the "only if" part of Theorem 4.5.1. $\square$

In order to prove the "if" part of Theorem 4.5.1, we recall from [Kam1, §2.3] the notion of Berenstein-Zelevinsky datum. We set $\Gamma := \{w \cdot \Lambda_i \mid w \in W,\, i \in I\}$, where Λ_i, $i \in I$, are the fundamental weights for $\mathfrak{g}$. Let $P = P(\mu_\bullet) \in \mathcal{MV}(\lambda)$ be an MV polytope with GGMS datum $\mu_\bullet = (\mu_w)_{w \in W}$. For each $\gamma \in \Gamma$, we define $M_\gamma \in \mathbb{Q}$ by:

$$M_\gamma := \langle w \cdot \Lambda_i,\, \mu_w \rangle \in \mathbb{Q} \quad \text{if } \gamma = w \cdot \Lambda_i \text{ for some } w \in W \text{ and } i \in I;$$

note that the rational number $M_\gamma \in \mathbb{Q}$ does note depend on the expression $\gamma = w \cdot \Lambda_i$, $w \in W$, $i \in I$, of $\gamma \in \Gamma$. We call the collection $M_\bullet = (M_\gamma)_{\gamma \in \Gamma}$ of rational numbers the Berenstein-Zelevinsky (BZ for short) datum of the MV polytope P. We know from [Kam1, Proposition 2.2] that

$$P = P(\mu_\bullet) = \{h \in \mathfrak{h}_\mathbb{R} \mid \langle \gamma,\, h \rangle \ge M_\gamma \text{ for all } \gamma \in \Gamma\}. \tag{4.5.2}$$

Lemma 4.5.4. *Let* $P_1 = P(\mu_\bullet^{(1)})$, $P_2 = P(\mu_\bullet^{(2)}) \in \mathcal{MV}(\lambda)$ *be MV polytopes with GGMS data* $\mu_\bullet^{(1)} = (\mu_w^{(1)})_{w \in W}$, $\mu_\bullet^{(2)} = (\mu_w^{(2)})_{w \in W}$, *and denote by* $M_\bullet^{(1)} = (M_\gamma^{(1)})_{\gamma \in \Gamma}$, $M_\bullet^{(2)} = (M_\gamma^{(2)})_{\gamma \in \Gamma}$ *the BZ data of* P_1, P_2, *respectively. Then,* $P_1 \subset P_2$ *if and only if* $M_\gamma^{(1)} \ge M_\gamma^{(2)}$ *for all* $\gamma \in \Gamma$.

Proof. The "if" part is obvious by (4.5.2). We show the "only if" part. Since $P_1 = \operatorname{Conv}\{\mu_w^{(1)} \mid w \in W\}$, and since $P_1 \subset P_2$ by assumption, it follows that $\mu_w^{(1)} \in P_2$ for all $w \in W$. Take an arbitrary element $\gamma = w \cdot \Lambda_i$, $w \in W$, $i \in I$, of Γ. Because $\mu_w^{(1)} \in P_2$ as above, we have $\langle \gamma,\, \mu_w^{(1)} \rangle \ge M_\gamma^{(2)}$ by (4.5.2). Hence, by the definition of BZ data, we get $M_\gamma^{(1)} = \langle w \cdot \Lambda_i,\, \mu_w^{(1)} \rangle = \langle \gamma,\, \mu_w^{(1)} \rangle \ge M_\gamma^{(2)}$. This proves the lemma. $\square$

Proof of the "if" part of Theorem 4.5.1. Let $P = P(\mu_\bullet) \in \mathcal{MV}(\lambda)$ be an MV polytope with GGMS datum $\mu_\bullet = (\mu_w)_{w \in W}$ such that $P \supset P_{x \cdot \lambda}$, and denote by $M_\bullet = (M_\gamma)_{\gamma \in \Gamma}$ the BZ datum of P. Take $\mathbf{i} = (i_1, i_2, \dots, i_m) \in R(w_0)$ such that $w_p^{\mathbf{i}} = s_{i_1} s_{i_2} \cdots s_{i_p} = xw_0$, with $p = \ell(xw_0)$. In order to prove the "if" part, it suffices to show that $M_{w_{l-1}^{\mathbf{i}} \cdot \Lambda_{i_l}} = M_{w_l^{\mathbf{i}} \cdot \Lambda_{i_l}}$ for all $p+1 \le l \le m$. Indeed, we know from [Kam1, Theorem 6.6] that for $1 \le l \le m$,

$$N_l := \varphi_{i_l}(f_{i_{l-1}}^{\max} \cdots f_{i_2}^{\max} f_{i_1}^{\max} P)$$

is equal to $M_{w_{l-1}^{\mathbf{i}} \cdot \Lambda_{i_l}} - M_{w_l^{\mathbf{i}} \cdot \Lambda_{i_l}}$. Hence, if $M_{w_{l-1}^{\mathbf{i}} \cdot \Lambda_{i_l}} = M_{w_l^{\mathbf{i}} \cdot \Lambda_{i_l}}$ holds for all $p+1 \le l \le m$, then it follows that $N_l = 0$ for all $p+1 \le l \le m$. Therefore, the argument after Claim 2 in the proof of Theorem 3.5.1 shows that $P \in \Psi_\lambda(\mathcal{B}^x(\lambda)) = \mathcal{MV}^x(\lambda)$.

We will show that $M_{w_{l-1}^{\mathbf{i}} \cdot \Lambda_{i_l}} = M_{w_l^{\mathbf{i}} \cdot \Lambda_{i_l}}$ for all $p+1 \le l \le m$. By the definition of $\mathcal{MV}(\lambda)$, the polytope $P \in \mathcal{MV}(\lambda)$ is contained in $\operatorname{Conv}(W \cdot \lambda) = P_{w_0 \cdot \lambda}$. Hence we have

$$P_{x \cdot \lambda} \subset P \subset \operatorname{Conv}(W \cdot \lambda) = P_{w_0 \cdot \lambda}.$$

Therefore, if we denote by $M_\bullet^{(1)} = (M_\gamma^{(1)})_{\gamma\in\Gamma}$ and $M_\bullet^{(2)} = (M_\gamma^{(2)})_{\gamma\in\Gamma}$ the BZ data of $P_{x\cdot\lambda}$ and $P_{w_0\cdot\lambda}$, respectively, we see from Lemma 4.5.4 that

$$M_\gamma^{(1)} \geq M_\gamma \geq M_\gamma^{(2)} \quad \text{for all } \gamma \in \Gamma. \tag{4.5.3}$$

Claim. *For all $p+1 \leq l \leq m$, we have*

$$M^{(1)}_{w^{\mathbf{i}}_{l-1}\cdot\Lambda_{i_l}} = M^{(2)}_{w^{\mathbf{i}}_{l-1}\cdot\Lambda_{i_l}} = \langle \Lambda_{i_l},\, w_0\cdot\lambda\rangle, \qquad M^{(1)}_{w^{\mathbf{i}}_{l}\cdot\Lambda_{i_l}} = M^{(2)}_{w^{\mathbf{i}}_{l}\cdot\Lambda_{i_l}} = \langle \Lambda_{i_l},\, w_0\cdot\lambda\rangle.$$

Proof of Claim. Let $\mu_\bullet^{(1)} = (\mu_w^{(1)})_{w\in W}$ and $\mu_\bullet^{(2)} = (\mu_w^{(2)})_{w\in W}$ be the GGMS data of $P_{x\cdot\lambda}$ and $P_{w_0\cdot\lambda}$, respectively. We deduce (see Example 4.1.2) that $\mu^{(1)}_{w^{\mathbf{i}}_l} = w^{\mathbf{i}}_l w_0\cdot\lambda$ for all $p \leq l \leq m$. Therefore, we have

$$\begin{aligned} M^{(1)}_{w^{\mathbf{i}}_{l-1}\cdot\Lambda_{i_l}} &= \langle w^{\mathbf{i}}_{l-1}\cdot\Lambda_{i_l},\, \mu_{w^{\mathbf{i}}_{l-1}}\rangle = \langle w^{\mathbf{i}}_{l-1}\cdot\Lambda_{i_l},\, w^{\mathbf{i}}_{l-1}w_0\cdot\lambda\rangle = \langle\Lambda_{i_l},\, w_0\cdot\lambda\rangle, \\ M^{(1)}_{w^{\mathbf{i}}_{l}\cdot\Lambda_{i_l}} &= \langle w^{\mathbf{i}}_{l}\cdot\Lambda_{i_l},\, \mu_{w^{\mathbf{i}}_{l}}\rangle = \langle w^{\mathbf{i}}_{l}\cdot\Lambda_{i_l},\, w^{\mathbf{i}}_{l}w_0\cdot\lambda\rangle = \langle\Lambda_{i_l},\, w_0\cdot\lambda\rangle. \end{aligned}$$

Also, we know (see Remark 2.2.4 (2)) that $\mu^{(2)}_{w^{\mathbf{i}}_l} = w^{\mathbf{i}}_l w_0\cdot\lambda$ for all $0 \leq l \leq m$. Hence, by the same calculation as above, we get

$$M^{(2)}_{w^{\mathbf{i}}_{l-1}\cdot\Lambda_{i_l}} = \langle\Lambda_{i_l},\, w_0\cdot\lambda\rangle, \qquad M^{(2)}_{w^{\mathbf{i}}_{l}\cdot\Lambda_{i_l}} = \langle\Lambda_{i_l},\, w_0\cdot\lambda\rangle.$$

This proves the claim. ▮

Let us take $p+1 \leq l \leq m$ arbitrarily. By combining the claim above with (4.5.3), we see that

$$\langle\Lambda_{i_l},\, w_0\cdot\lambda\rangle = M^{(1)}_{w^{\mathbf{i}}_{l-1}\cdot\Lambda_{i_l}} \geq M_{w^{\mathbf{i}}_{l-1}\cdot\Lambda_{i_l}} \geq M^{(2)}_{w^{\mathbf{i}}_{l-1}\cdot\Lambda_{i_l}} = \langle\Lambda_{i_l},\, w_0\cdot\lambda\rangle,$$

and hence that $M_{w^{\mathbf{i}}_{l-1}\cdot\Lambda_{i_l}} = \langle\Lambda_{i_l},\, w_0\cdot\lambda\rangle$. Similarly, we see that $M_{w^{\mathbf{i}}_{l}\cdot\Lambda_{i_l}} = \langle\Lambda_{i_l},\, w_0\cdot\lambda\rangle$. Thus, we obtain $M_{w^{\mathbf{i}}_{l-1}\cdot\Lambda_{i_l}} = \langle\Lambda_{i_l},\, w_0\cdot\lambda\rangle = M_{w^{\mathbf{i}}_{l}\cdot\Lambda_{i_l}}$. This completes the proof of the "if" part of Theorem 4.5.1. ☐

4.6 Open problem.

In view of Theorem 4.5.1, it is natural to pose the following question.

Open Problem. Let us take an arbitrary $x \in W$. Are the MV polytopes lying in the Demazure crystal $\mathcal{MV}_x(\lambda)$ all contained (as a set) in the extremal MV polytope $P_{x\cdot\lambda} = \mathrm{Conv}(W_{\leq x}\cdot\lambda)$?

Remark 4.6.1. We cannot expect that the converse statement holds, as the following example shows. Let $\mathfrak{g}$ be the simple Lie algebra of type A_2; the set $R(w_0)$ consists of two elements $\mathbf{i} := (1,2,1)$ and $\mathbf{j} := (2,1,2)$. Set $x := s_1s_2 \in W$ and $\lambda := h_1 + h_2 \in X_*(T) \subset \mathfrak{h}_\mathbb{R}$. Then, the MV polytope $f_2f_1P_\lambda \in \mathcal{MV}(\lambda)$ is contained (as a set) in the extremal MV polytope $P_{x\cdot\lambda} = \mathrm{Conv}(W_{\leq x}\cdot\lambda)$, but $f_2f_1P_\lambda$ is not an element of the Demazure crystal $\mathcal{MV}_x(\lambda)$.

Indeed, let $\mu_\bullet = (\mu_w)_{w\in W}$ be the GGMS datum of the MV polytope $f_2 f_1 P_\lambda$. Using the definition of the Kashiwara operators f_1, f_2 and the formula (2.1.6), we deduce that the integers $n^{\mathbf{i}}_l = n^{\mathbf{i}}_l(\mu_\bullet)$, $1 \le l \le 3$, and $n^{\mathbf{j}}_l = n^{\mathbf{j}}_l(\mu_\bullet)$, $1 \le l \le 3$, are given as:

$$\begin{cases} n^{\mathbf{i}}_1 = 0, & n^{\mathbf{i}}_2 = 1, \qquad n^{\mathbf{i}}_3 = 0, \\ n^{\mathbf{j}}_1 = 1, & n^{\mathbf{j}}_2 = 0, \qquad n^{\mathbf{j}}_3 = 1, \end{cases} \tag{4.6.1}$$

and hence the GGMS datum $\mu_\bullet = (\mu_w)_{w\in W}$ of $f_2 f_1 P_\lambda$ is given as:

$$\begin{aligned} &\mu_{w_0} = \mu_{w^{\mathbf{i}}_3} = \mu_{w^{\mathbf{j}}_3} = \lambda, \qquad \mu_{s_1 s_2} = \mu_{w^{\mathbf{i}}_2} = \lambda, \\ &\mu_{s_1} = \mu_{w^{\mathbf{i}}_1} = \lambda - (h_1 + h_2) = 0, \qquad \mu_e = \mu_{w^{\mathbf{i}}_0} = \mu_{w^{\mathbf{j}}_0} = \lambda - (h_1 + h_2) = 0, \\ &\mu_{s_2 s_1} = \mu_{w^{\mathbf{j}}_2} = \lambda - h_1 = s_1 \cdot \lambda, \qquad \mu_{s_2} = \mu_{w^{\mathbf{j}}_1} = \lambda - h_1 = s_1 \cdot \lambda. \end{aligned}$$

From these, it is easily seen that the MV polytope $f_2 f_1 P_\lambda$ is contained in the extremal MV polytope $P_{x\cdot\lambda} = \mathrm{Conv}(W_{\le x}\cdot\lambda)$ (see the figures below). However, since $x w_0 = s_1 s_2 s_2 s_1 s_2 = s_2$, and hence $S(xw_0, \mathbf{i}) = \{(2)\}$ and $S(xw_0, \mathbf{j}) = \{(1), (3)\}$, it follows from (4.6.1) that $f_2 f_1 P_\lambda$ is not an element of $\mathcal{MV}_x(\lambda)$.





Remark 4.6.2. We know (see [BaG, §4.1] and [Kam2, Theorem 4.7]) that for $P \in \mathcal{MV}(\lambda)$ and $j \in I$ with $e_j P \ne \mathbf{0}$, there holds $e_j P \subset P$. Therefore, if an MV polytope $P \in \mathcal{MV}_x(\lambda)$ were obtained from the extremal MV polytope $P_{x\cdot\lambda}$ by successive application of the Kashiwara operators e_j, $j \in I$, then it would follow immediately that the polytope P is contained in $P_{x\cdot\lambda}$. However, in general, not all MV polytopes in $\mathcal{MV}_x(\lambda)$ can be obtained from $P_{x\cdot\lambda}$ by successively applying e_j, $j \in I$, as Example 4.6.3 below shows.

Example 4.6.3. As in Remark 4.6.1, let $\mathfrak{g}$ be the simple Lie algebra of type A_2, and set $x = s_1 s_2 \in W$ and $\lambda = h_1 + h_2 \in X_*(T) \subset \mathfrak{h}_\mathbb{R}$; the set $R(w_0)$ consists of $\mathbf{i} = (1, 2, 1)$ and $\mathbf{j} = (2, 1, 2)$. Then, $f_1 P_\lambda \in \mathcal{MV}_x(\lambda)$. Indeed, let $\mu_\bullet = (\mu_w)_{w\in W}$ be the GGMS datum of the MV polytope $f_1 P_\lambda$. Using the definition of the Kashiwara operator f_1 and the formula (2.1.6), we deduce that the integers $n^{\mathbf{i}}_l = n^{\mathbf{i}}_l(\mu_\bullet)$, $1 \le l \le 3$, and $n^{\mathbf{j}}_l = n^{\mathbf{j}}_l(\mu_\bullet)$, $1 \le l \le 3$, are given as:

$$\begin{cases} n^{\mathbf{i}}_1 = 1, & n^{\mathbf{i}}_2 = 0, \qquad n^{\mathbf{i}}_3 = 0, \\ n^{\mathbf{j}}_1 = 0, & n^{\mathbf{j}}_2 = 0, \qquad n^{\mathbf{j}}_3 = 1. \end{cases} \tag{4.6.2}$$

Since $xw_0 = s_1s_2s_2s_1s_2 = s_2$, and hence $S(xw_0, \mathbf{i}) = \{(2)\}$ and $S(xw_0, \mathbf{j}) = \{(1), (3)\}$, it follows from (4.6.2) that f_1P_λ is an element of $\mathcal{MV}_x(\lambda)$. However, we deduce from the crystal graph (4.6.3) below of $\mathcal{MV}(\lambda) \cong \mathcal{B}(\lambda)$ that f_1P_λ cannot be obtained from $P_{x\cdot\lambda}$ by successively applying e_1 and e_2.



(4.6.3)

5 Decomposition of Demazure crystals and opposite Demazure crystals.

We fix (once and for all) an arbitrary dominant coweight $\lambda \in X_*(T) \subset \mathfrak{h}_\mathbb{R}$.

5.1 Decomposition of Demazure crystals. We set $W_\lambda := \{w \in W \mid w \cdot \lambda = \lambda\} \subset W$, and denote by $W^\lambda_{\min}$ the set of minimal coset representatives for the quotient set W/W_λ with respect to the Bruhat ordering $\leq$ on W (see, for example, [BjB, Chap. 2, §4]). For each $x \in W^\lambda_{\min}$, define a subset $\overline{\mathcal{MV}}_x(\lambda)$ of the Demazure crystal $\mathcal{MV}_x(\lambda)$ by (cf. [Kas4, §9.1]):

$$\overline{\mathcal{MV}}_x(\lambda) = \mathcal{MV}_x(\lambda) \setminus \left(\bigcup_{z \in W^\lambda_{\min},\ z<x} \mathcal{MV}_z(\lambda) \right); \tag{5.1.1}$$

recall from [Kas2, Proposition 3.2.4] that for $z \in W^\lambda_{\min}$ with $z < x$, we have $\mathcal{MV}_z(\lambda) \subset \mathcal{MV}_x(\lambda)$. We can easily show by induction on $\ell(x)$ (see also [Kas4, §9.1]) that for each $x \in W$,

$$\mathcal{MV}_x(\lambda) = \bigsqcup_{z \in W^\lambda_{\min},\ z \leq x} \overline{\mathcal{MV}}_z(\lambda), \tag{5.1.2}$$

and in particular,

$$\mathcal{MV}(\lambda) = \bigsqcup_{z \in W^\lambda_{\min}} \overline{\mathcal{MV}}_z(\lambda); \tag{5.1.3}$$

note that $\mathcal{MV}_{w_0}(\lambda) = \mathcal{MV}(\lambda)$. Similarly, for each $x \in W$, define a subset $\overline{\mathcal{MV}}_x(\infty)$ of the Demazure crystal $\mathcal{MV}_x(\infty)$ by (cf. [Kas4, Proposition 9.1.6 (2)]):

$$\overline{\mathcal{MV}}_x(\infty) = \mathcal{MV}_x(\infty) \setminus \left(\bigcup_{z \in W,\ z < x} \mathcal{MV}_z(\infty) \right).$$

It follows from [Kas4, Proposition 9.1.6 (2)] that for each $x \in W$,

$$\mathcal{MV}_x(\infty) = \bigsqcup_{z \in W,\ z \leq x} \overline{\mathcal{MV}}_z(\infty), \tag{5.1.4}$$

and in particular,

$$\mathcal{MV}(\infty) = \bigsqcup_{z \in W} \overline{\mathcal{MV}}_z(\infty); \tag{5.1.5}$$

note that $\mathcal{MV}_{w_0}(\infty) = \mathcal{MV}(\infty)$. For each $P \in \mathcal{MV}(\lambda)$ (resp., $P \in \mathcal{MV}(\infty)$), we denote by $\iota(P)$ the unique element $z \in W^{\lambda}_{\mathrm{min}}$ (resp., $z \in W$) for which $P \in \overline{\mathcal{MV}}_z(\lambda)$ (resp., $P \in \overline{\mathcal{MV}}_z(\infty)$); we see from (5.1.2) and (5.1.3) (resp., (5.1.4) and (5.1.5)) that the element $\iota(P) \in W$ specifies the smallest (with respect to the inclusion relation) Demazure crystal in which P lies.

Remark 5.1.1. Let $x \in W$, and denote by $x^{\lambda}_{\mathrm{min}} \in W^{\lambda}_{\mathrm{min}}$ the (unique) minimal element of the left coset xW_{λ} with respect to the Bruhat ordering on W. From (5.1.1), it is easily seen that $\iota(P_{x \cdot \lambda}) = x^{\lambda}_{\mathrm{min}}$.

Let $P = P(\mu_{\bullet})$ be an element of $\mathcal{MV}(\lambda)$ (resp., $\mathcal{MV}(\infty)$) with GGMS datum $\mu_{\bullet} = (\mu_w)_{w \in W}$. For each $\mathbf{i} = (i_1, i_2, \ldots, i_m) \in R(w_0)$, we set

$$T(P, \mathbf{i}) := \left\{ (a_1, a_2, \ldots, a_l) \in [1, m]^l \;\middle|\; \begin{array}{l} 0 \leq l \leq m \\ 1 \leq a_1 < a_2 < \cdots < a_l \leq m \\ n^{\mathbf{i}}_{a_q} = n^{\mathbf{i}}_{a_q}(\mu_{\bullet}) = 0 \quad \text{for all } 1 \leq q \leq l \end{array} \right\},$$

and define a subset $W(P, \mathbf{i})$ of W by:

$$W(P, \mathbf{i}) = \bigl\{ s_{i_{a_1}} s_{i_{a_2}} \cdots s_{i_{a_l}} w_0 \in W \mid (a_1, a_2, \ldots, a_l) \in T(P, \mathbf{i}) \bigr\}.$$

Lemma 5.1.2. *Let $P \in \mathcal{MV}(\lambda)$ (resp., $P \in \mathcal{MV}(\infty)$), and take an arbitrary $\mathbf{i} \in R(w_0)$. We have $\iota(P) \in W(P, \mathbf{i})$, and $z \geq \iota(P)$ for all $z \in W(P, \mathbf{i})$ with respect to the Bruhat ordering $\geq$ on W.*

Proof. We give a proof only for the case that $P \in \mathcal{MV}(\lambda)$; the proof for the case that $P \in \mathcal{MV}(\infty)$ is similar. Since $P \in \overline{\mathcal{MV}}_{\iota(P)}(\lambda) \subset \mathcal{MV}_{\iota(P)}(\lambda)$ by the definition of $\iota(P)$, it follows from Corollary 3.2.2 that there exists a sequence $(a_1, a_2, \ldots, a_l) \in \widehat{S}(\iota(P)w_0, \mathbf{i})$ such that $n^{\mathbf{i}}_{a_q} = n^{\mathbf{i}}_{a_q}(\mu_{\bullet}) = 0$ for all $1 \leq q \leq l$. Therefore, the sequence $(a_1, a_2, \ldots, a_l)$ is an element of $T(P, \mathbf{i})$, and hence

$$\iota(P) = (\iota(P)w_0)w_0 = s_{i_{a_1}} s_{i_{a_2}} \cdots s_{i_{a_l}} w_0 \in W(P, \mathbf{i}).$$

Now, let $z \in W(P, \mathbf{i})$. Then we deduce from Corollary 3.2.2 that $P \in \mathcal{MV}_z(\lambda)$. Denote by $z_{\min}^{\lambda} \in W_{\min}^{\lambda}$ the minimal element of the left coset zW_λ; obviously, $z_{\min}^{\lambda} \le z$. Since $z_{\min}^{\lambda} \cdot \lambda = z \cdot \lambda$, we have $P \in \mathcal{MV}_z(\lambda) = \mathcal{MV}_{z_{\min}^{\lambda}}(\lambda)$ (see Remark 3.1.1). Therefore, by the decomposition (5.1.2), there exists an element $z' \in W_{\min}^{\lambda}$ with $z' \le z_{\min}^{\lambda}$ such that $P \in \overline{\mathcal{MV}}_{z'}(\lambda)$. It follows from the uniqueness of $\iota(P)$ that $z' = \iota(P)$, and hence $\iota(P) \le z_{\min}^{\lambda}$. Since $z_{\min}^{\lambda} \le z$ as noted above, we get $\iota(P) \le z$. This proves the lemma. □

Remark 5.1.3. Let $P \in \mathcal{MV}(\lambda)$ (resp., $P \in \mathcal{MV}(\infty)$), and take an arbitrary $\mathbf{i} \in R(w_0)$. By Lemma 5.1.2, $\iota(P)$ is the minimum element $\min W(P, \mathbf{i})$ of $W(P, \mathbf{i})$ with respect to the Bruhat ordering on W. Namely, we have $\iota(P) = \min W(P, \mathbf{i})$ for every $\mathbf{i} \in R(w_0)$.

The next proposition follows immediately from Remark 5.1.3.

Proposition 5.1.4. *Let $P \in \mathcal{MV}(\lambda)$ (resp., $P \in \mathcal{MV}(\infty)$), and let $x \in W_{\min}^{\lambda}$ (resp., $x \in W$). Then, $P \in \overline{\mathcal{MV}}_x(\lambda)$ (resp., $P \in \overline{\mathcal{MV}}_x(\infty)$) if and only if for some (or equivalently, every) $\mathbf{i} \in R(w_0)$, the element x is identical to the minimum element $\min W(P, \mathbf{i})$ of $W(P, \mathbf{i})$ with respect to the Bruhat ordering on W.*

Here we give an application of our description above for the elements $\iota(P)$, $P \in \mathcal{MV}(\lambda)$. Let $* : U_q^-(\mathfrak{g}^\vee) \to U_q^-(\mathfrak{g}^\vee)$ be the $\mathbb{C}(q)$-algebra antiautomorphism that fixes the Chevalley generators $y_j \in U_q^-(\mathfrak{g}^\vee)$, $j \in I$, corresponding to the negative simple roots (see [Kas1, §1.3]). We know from [Kas2, Theorem 2.1.1] that this antiautomorphism $* : U_q^-(\mathfrak{g}^\vee) \to U_q^-(\mathfrak{g}^\vee)$ induces an involution $* : \mathcal{B}(\infty) \to \mathcal{B}(\infty)$ on the crystal base $\mathcal{B}(\infty)$ of $U_q^-(\mathfrak{g}^\vee)$, which we call the Kashiwara involution (or, $*$-operation) on $\mathcal{B}(\infty)$. We then define an involution $* : \mathcal{MV}(\infty) \to \mathcal{MV}(\infty)$ so that the following diagram commutes:

$$\begin{tikzcd}
\mathcal{MV}(\infty) \arrow[r, "*"] \arrow[d, "\Psi"'] & \mathcal{MV}(\infty) \arrow[d, "\Psi"] \\
\mathcal{B}(\infty) \arrow[r, "*"] & \mathcal{B}(\infty),
\end{tikzcd}$$

and for $P \in \mathcal{MV}(\infty)$, denote by P^* the image of P under the involution $* : \mathcal{MV}(\infty) \to \mathcal{MV}(\infty)$.

Corollary 5.1.5. *Let $P \in \mathcal{MV}(\infty)$. Then, the element $\iota(P^*) \in W$ is identical to the inverse $\bigl(\iota(P)\bigr)^{-1}$ of the element $\iota(P) \in W$.*

Proof of Corollary 5.1.5. Let $\mu_\bullet = (\mu_w)_{w \in W}$ and $\mu'_\bullet = (\mu'_w)_{w \in W}$ be GGMS data for $P \in \mathcal{MV}(\infty)$ and $P^* \in \mathcal{MV}(\infty)$, respectively. Take an arbitrary $\mathbf{i} = (i_1, i_2, \ldots, i_m) \in R(w_0)$, and define $\mathbf{j} = (j_1, j_2, \ldots, j_m) \in R(w_0)$ by: $\mathbf{j} = (\omega(i_m), \omega(i_{m-1}), \ldots, \omega(i_1))$, where $\omega : I \to I$ is the Dynkin diagram automorphism for $\mathfrak{g}$ such that $\alpha_{\omega(j)} = -w_0 \cdot \alpha_j$ for all $j \in I$. Then we deduce from [Kam2, Proposition 6.1] (see also [BeZ, Proposition 3.3 (iii)]) that

$$\Bigl(n_1^{\mathbf{j}}(\mu'_\bullet),\, n_2^{\mathbf{j}}(\mu'_\bullet),\, \ldots,\, n_m^{\mathbf{j}}(\mu'_\bullet)\Bigr) = \Bigl(n_m^{\mathbf{i}}(\mu_\bullet),\, n_{m-1}^{\mathbf{i}}(\mu_\bullet),\, \ldots,\, n_1^{\mathbf{i}}(\mu_\bullet)\Bigr). \tag{5.1.6}$$

From (5.1.6), it follows that a sequence $(a_1, a_2, \dots, a_l)$ is an element of $T(P, \mathbf{i})$ if and only if the sequence $(m - a_l + 1, m - a_{l-1} + 1, \dots, m - a_1 + 1)$ is an element of $T(P^*, \mathbf{j})$. Therefore, we have

$$\begin{aligned} W(P^*, \mathbf{j}) &= \bigl\{ s_{j_{b_1}} s_{j_{b_2}} \cdots s_{j_{b_l}} w_0 \in W \mid (b_1, b_2, \dots, b_l) \in T(P^*, \mathbf{j}) \bigr\} \\ &= \bigl\{ s_{j_{m-a_l+1}} s_{j_{m-a_{l-1}+1}} \cdots s_{j_{m-a_1+1}} w_0 \in W \mid (a_1, a_2, \dots, a_l) \in T(P, \mathbf{i}) \bigr\} \\ &= \bigl\{ s_{i_{\omega(a_l)}} s_{i_{\omega(a_{l-1})}} \cdots s_{i_{\omega(a_1)}} w_0 \in W \mid (a_1, a_2, \dots, a_l) \in T(P, \mathbf{i}) \bigr\} \\ &= \bigl\{ w_0 s_{i_{a_l}} s_{i_{a_{l-1}}} \cdots s_{i_{a_1}} \in W \mid (a_1, a_2, \dots, a_l) \in T(P, \mathbf{i}) \bigr\} \\ &= \bigl\{ w^{-1} \mid w \in W(P, \mathbf{i}) \bigr\}. \end{aligned} \tag{5.1.7}$$

We know from Proposition 5.1.4 that the elements $\iota(P^*)$ and $\iota(P)$ are identical to the minimum elements $\min W(P^*, \mathbf{j})$ and $\min W(P, \mathbf{i})$, respectively. Combining this fact and (5.1.7) above, we conclude that $\iota(P^*) = \bigl(\iota(P)\bigr)^{-1}$, as desired. ☐

5.2 Decomposition of opposite Demazure crystals. Let us denote by $W^{\lambda}_{\max}$ the set of maximal coset representatives for the quotient set W/W_λ with respect to the Bruhat ordering $\le$ on W (see, for example, [BjB, Chap. 2, §4]).

Remark 5.2.1. It is clear that $W^{\lambda}_{\max} = W^{\lambda}_{\min} w_{0,\lambda}$, where $w_{0,\lambda} \in W_\lambda$ denotes the longest element of W_λ.

For each $x \in W^{\lambda}_{\max}$, define a subset $\overline{\mathcal{MV}}^{x}(\lambda)$ of the opposite Demazure crystal $\mathcal{MV}^{x}(\lambda)$ by:

$$\overline{\mathcal{MV}}^{x}(\lambda) = \mathcal{MV}^{x}(\lambda) \setminus \left(\bigcup_{z \in W^{\lambda}_{\max},\ z > x} \mathcal{MV}^{z}(\lambda) \right); \tag{5.2.1}$$

recall from [Kas2, Proposition 3.2.4] (along with the comment following [Kas2, Proposition 4.3]) that for $z \in W^{\lambda}_{\max}$ with $z > x$, we have $\mathcal{MV}^{z}(\lambda) \subset \mathcal{MV}^{x}(\lambda)$. We can easily show by descending induction on $\ell(x)$ that for each $x \in W$,

$$\mathcal{MV}^{x}(\lambda) = \bigsqcup_{z \in W^{\lambda}_{\max},\ z \ge x} \overline{\mathcal{MV}}^{z}(\lambda), \tag{5.2.2}$$

and in particular,

$$\mathcal{MV}(\lambda) = \bigsqcup_{z \in W^{\lambda}_{\max}} \overline{\mathcal{MV}}^{z}(\lambda); \tag{5.2.3}$$

note that $\mathcal{MV}^{e}(\lambda) = \mathcal{MV}(\lambda)$. For each $P \in \mathcal{MV}(\lambda)$, we denote by $\kappa(P)$ the unique element $z \in W^{\lambda}_{\max}$ for which $P \in \overline{\mathcal{MV}}^{z}(\lambda)$; we see from (5.2.2) and (5.2.3) that the element $\kappa(P) \in W$ specifies the smallest (with respect to the inclusion relation) opposite Demazure crystal in which P lies.

Remark 5.2.2. Let $x \in W$, and denote by $x^{\lambda}_{\max} \in W^{\lambda}_{\max}$ the (unique) maximal element of the left coset xW_λ with respect to the Bruhat ordering on W; note that $x^{\lambda}_{\max} = x^{\lambda}_{\min} w_{0,\lambda}$. From (5.2.1), it is easily seen that $\kappa(P_{x\cdot\lambda}) = x^{\lambda}_{\max} = x^{\lambda}_{\min} w_{\lambda,0}$.

Let $P = P(\mu_\bullet)$ be an element of $\mathcal{MV}(\lambda)$ with GGMS datum $\mu_\bullet = (\mu_w)_{w \in W}$. For each $\mathbf{i} = (i_1, i_2, \ldots, i_m) \in R(w_0)$, we set

$$U(P, \mathbf{i}) := \bigl\{ 0 \le p \le m \mid \mu_{w_l^{\mathbf{i}}} = w_l^{\mathbf{i}} w_0 \cdot \lambda \text{ for every } p \le l \le m \bigr\}.$$

Then we set

$$Y(P) := \bigl\{ w_p^{\mathbf{i}} w_0 \mid p \in U(P, \mathbf{i}),\, \mathbf{i} \in R(w_0) \bigr\}.$$

Lemma 5.2.3. *Let $P \in \mathcal{MV}(\lambda)$. We have $\kappa(P) \in Y(P)$, and $\kappa(P) \ge z$ for all $z \in Y(P)$ with respect to the Bruhat ordering $\ge$ on W.*

Proof. Let $P = P(\mu_\bullet) \in \mathcal{MV}(\lambda)$ be an MV polytope with GGMS datum $\mu_\bullet = (\mu_w)_{w \in W}$. It follows from the definition of $\kappa(P)$ that $P \in \overline{\mathcal{MV}}^{\kappa(P)}(\lambda) \subset \mathcal{MV}^{\kappa(P)}(\lambda)$. Therefore, by the definition of $\mathcal{MV}^{\kappa(P)}(\lambda)$, there exists $\mathbf{i} \in R(w_0)$ such that $w_p^{\mathbf{i}} = \kappa(P) w_0$, with $p = \ell(\kappa(P) w_0)$, and such that $\mu_{w_l^{\mathbf{i}}} = w_l^{\mathbf{i}} w_0 \cdot \lambda$ for all $p \le l \le m$. Hence we have

$$Y(P) \ni w_p^{\mathbf{i}} w_0 = \bigl(\kappa(P) w_0\bigr) w_0 = \kappa(P).$$

Now, let $z \in Y(P)$. Then we deduce from Theorem 3.5.1 that $P \in \mathcal{MV}^z(\lambda)$. Denote by $z_{\max}^\lambda \in W_{\max}^\lambda$ the maximal element of the left coset zW_λ; obviously, $z_{\max}^\lambda \ge z$. Since $z_{\max}^\lambda \cdot \lambda = z \cdot \lambda$, we have $P \in \mathcal{MV}^z(\lambda) = \mathcal{MV}^{z_{\max}^\lambda}(\lambda)$ (see Remark 3.4.1). Therefore, by the decomposition (5.2.2), there exists an element $z' \in W_{\max}^\lambda$ with $z' \ge z_{\max}^\lambda$ such that $P \in \overline{\mathcal{MV}}^{z'}(\lambda)$. It follows from the uniqueness of $\kappa(P)$ that $z' = \kappa(P)$, and hence $\kappa(P) \ge z_{\max}^\lambda$. Since $z_{\max}^\lambda \ge z$ as noted above, we get $\kappa(P) \ge z$. This proves the lemma. □

Remark 5.2.4. Let $P \in \mathcal{MV}(\lambda)$. By Lemma 5.2.3, $\kappa(P)$ is the maximum element $\max Y(P)$ of $Y(P)$ with respect to the Bruhat ordering on W. Namely, we have $\kappa(P) = \max Y(P)$.

The next proposition follows immediately from Remark 5.2.4.

Proposition 5.2.5. *Let $P \in \mathcal{MV}(\lambda)$, and let $x \in W_{\max}^\lambda$. Then, $P \in \overline{\mathcal{MV}}^x(\lambda)$ if and only if the element x is identical to the maximum element $\max Y(P)$ of $Y(P)$ with respect to the Bruhat ordering on W.*

References

[An] J. E. Anderson, A polytope calculus for semisimple groups, *Duke Math. J.* **116** (2003), 567–588.

[At] M. F. Atiyah, Convexity and commuting Hamiltonians, *Bull. London Math. Soc.* **14** (1982), 1–15.

[AP] M. Atiyah and A. Pressley, Convexity and loop groups, *in* “Arithmetic and geometry, Vol. II” (M. Artin and J. Tate, Eds.), pp. 33–63, Progr. Math. Vol. 36, Birkhäuser, Boston, 1983.

[BaG] P. Baumann and S. Gaussent, On Mirković-Vilonen cycles and crystal combinatorics, *Represent. Theory* **12** (2008), 83–130.

[BeZ] A. Berenstein and A. Zelevinsky, Tensor product multiplicities, canonical bases and totally positive varieties, *Invent. Math.* **143** (2001), 77–128.

[BjB] A. Björner and F. Brenti, "Combinatorics of Coxeter Groups", Graduate Texts in Mathematics Vol. 231, Springer, New York, 2005.

[BFG] A. Braverman, M. Finkelberg, and D. Gaitsgory, Uhlenbeck spaces via affine Lie algebras, *in* "The Unity of Mathematics" (P. Etingof et al., Eds.), pp. 17–135, Progr. Math. Vol. 244, Birkhäuser, Boston, 2006.

[BrG] A. Braverman and D. Gaitsgory, Crystals via the affine Grassmannian, *Duke Math. J.* **107** (2001), 561–575.

[GL] S. Gaussent and P. Littelmann, LS galleries, the path model, and MV cycles, *Duke Math. J.* **127** (2005), 35–88.

[Ha] T.J. Haines, Equidimensionality of convolution morphisms and applications to saturation problems, *Adv. Math.* **207** (2006), 297–327.

[Ho] J. Hong, The action of a Dynkin automorphism on Mirković-Vilonen cycles and polytopes, preprint, 2007, arXiv:0711.0070.

[I] B. Ion, A weight multiplicity formula for Demazure modules, *Int. Math. Res. Not.* **2005** (2005), no. 5, 311–323.

[Kam1] J. Kamnitzer, Mirković-Vilonen cycles and polytopes, preprint, 2005, arXiv:math.AG/ 0501365.

[Kam2] J. Kamnitzer, The crystal structure on the set of Mirković-Vilonen polytopes, *Adv. Math.* **215** (2007), 66–93.

[Kas1] M. Kashiwara, On crystal bases of the q-analogue of universal enveloping algebras, *Duke Math. J.* **63** (1991), 465–516.

[Kas2] M. Kashiwara, The crystal base and Littelmann's refined Demazure character formula, *Duke Math. J.* **71** (1993), 839–858.

[Kas3] M. Kashiwara, On level-zero representations of quantized affine algebras, *Duke Math. J.* **112** (2002), 117–175.

[Kas4] M. Kashiwara, Bases cristallines des groupes quantiques, Cours Spécialisés Vol. 9, Société Mathématique de France, Paris, 2002.

[Kat] S. Kato, Private communication, 2008.

[LP] C. Lenart and A. Postnikov, Affine Weyl groups in K-Theory and representation theory, *Int. Math. Res. Not.* **2007** (2007), no. 12, Article ID rnm038, 65 pages.

[Li1] P. Littelmann, A Littlewood-Richardson rule for symmetrizable Kac-Moody algebra, *Invent. Math.* **116** (1994), 329–346.

[Li2] P. Littelmann, Crystal graphs and Young tableaux, *J. Algebra* **175** (1995), 65–87.

[Li3] P. Littelmann, Cones, crystals, and patterns, *Transform. Groups* **3** (1998), 145–179.

[Lu1] G. Lusztig, “Introduction to Quantum Groups”, Progr. Math. Vol. 110, Birkhäuser, Boston, 1993.

[Lu2] G. Lusztig, Braid group action and canonical bases, *Adv. Math.* **122** (1996), 237–261.

[MiV1] I. Mirković and K. Vilonen, Perverse sheaves on affine Grassmannians and Langlands duality, *Math. Res. Lett.* **7** (2000), 13–24.

[MiV2] I. Mirković and K. Vilonen, Geometric Langlands duality and representations of algebraic groups over commutative rings, *Ann. of Math.* (2) **166** (2007), 95–143.

[MoP] R. V. Moody and A. Pianzola, “Lie Algebras with Triangular Decompositions”, Canadian Mathematical Society Series of Monographs and Advanced Texts, Wiley-Interscience, New York, 1995.

[NS1] S. Naito and D. Sagaki, Lakshmibai-Seshadri paths fixed by a diagram automorphism, *J. Algebra* **245** (2001), 395–412.

[NS2] S. Naito and D. Sagaki, Crystal bases and diagram automorphisms, *in* “Representation Theory of Algebraic Groups and Quantum Groups” (T. Shoji et al., Eds.), pp. 321–341, Adv. Stud. Pure Math. Vol. 40, Math. Soc. Japan, Tokyo, 2004.

[NS3] S. Naito and D. Sagaki, A modification of the Anderson-Mirković conjecture for Mirković-Vilonen polytopes in types B and C, *J. Algebra* **320** (2008), 387–416.

[N] T. Nakashima, Polytopes for crystallized Demazure modules and extremal vectors, *Comm. Algebra* **30** (2002), 1349–1367.

[NP] B. C. Ngô and P. Polo, Résolutions de Demazure affines et formule de Casselman-Shalika géométrique, *J. Algebraic Geom.* **10** (2001), 515–547.

[Sa] A. Savage, Quiver varieties and Demazure modules, *Math. Ann.* **335** (2006), 31–46.

[Sc] C. Schwer, Galleries and q-analogs in combinatorial representation theory, Dissertation, University of Cologne, 2006.